

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number  
**WO 2004/029626 A1**

(51) International Patent Classification<sup>7</sup>: G01N 33/566,  
C07K 14/705, 14/72

SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:  
PCT/EP2003/010666

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date:  
25 September 2003 (25.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/413,840 27 September 2002 (27.09.2002) US  
60/442,891 28 January 2003 (28.01.2003) US

(71) Applicant (for all designated States except US): BAYER  
HEALTHCARE AG [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): INBE, Hisayo  
[JP/JP]; 1-8-3-404, Motomachi, Tomoi, Nara-shi, Nara  
631-0078 (JP). ENCINAS, Jeffrey [US/JP]; 464-1-1401,  
Toroyamacho, Nishikikoji-sagaru, Nishinotoindori, Nak-  
agyo-ku, Kyoto-shi, Kyoto 604-8225 (JP).

(74) Common Representative: BAYER HEALTHCARE  
AG; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH,  
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,  
TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO  
patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG,  
ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU,  
TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT,  
RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: REGULATION OF HUMAN P2Y15 G PROTEIN-COUPLED RECEPTOR

(57) Abstract: Reagents which regulate human P2Y15 G protein-coupled receptor can play a role in preventing, ameliorating,  
or correcting bronchoconstriction or inflammation in diseases such as allergies including but not limited to asthma. Further, such  
reagents is useful to treat diseases of kidney function or disease related to mast cell.



WO 2004/029626 A1

**REGULATION OF HUMAN P2Y15 G PROTEIN-COUPLED RECEPTOR****TECHNICAL FIELD OF THE INVENTION**

5

The invention relates to the area of G protein-coupled receptors. More particularly, it relates to the area of P2Y15 G protein-coupled receptors and their regulation. It further relates to the treatment of bronchoconstriction and inflammation.

10

**BACKGROUND OF THE INVENTION****G Protein-Coupled Receptors**

Many medically significant biological processes are mediated by signal transduction pathways that involve G proteins (Lefkowitz, *Nature* 351, 353-354, 1991). The family of G protein-coupled receptors (GPCR) includes receptors for hormones, neurotransmitters, growth factors, and viruses. Specific examples of GPCRs include receptors for such diverse agents as dopamine, calcitonin, adrenergic hormones, endothelin, cAMP, adenosine, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorants, cytomegalovirus, G proteins themselves, effector proteins such as phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins such as protein kinase A and protein kinase C.

20

25

30

GPCRs possess seven conserved membrane-spanning domains connecting at least eight divergent hydrophilic loops. GPCRs (also known as 7TM receptors) have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. Most GPCRs have single conserved cysteine residues in each of the first two extracellular loops, which form disulfide bonds that are believed to stabilize functional protein structure. The seven transmembrane regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 has been implicated in signal transduction.

- 2 -

Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues can influence signal transduction of some GPCRs. Most GPCRs contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus.

5 For several GPCRs, such as the  $\beta$ -adrenergic receptor, phosphorylation by protein kinase A and/or specific receptor kinases mediates receptor desensitization.

For some receptors, the ligand binding sites of GPCRs are believed to comprise hydrophilic sockets formed by several GPCR transmembrane domains. The hydro-  
10 philic sockets are surrounded by hydrophobic residues of the GPCRs. The hydrophilic side of each GPCR transmembrane helix is postulated to face inward and form a polar ligand binding site. TM3 has been implicated in several GPCRs as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine, and TM6 or TM7 phenylalanines or tyrosines also are implicated in  
15 ligand binding.

GPCRs are coupled inside the cell by heterotrimeric G-proteins to various intracellular enzymes, ion channels, and transporters (*see Johnson et al., Endoc. Rev. 10, 317-331, 1989*). Different G-protein alpha-subunits preferentially stimulate  
20 particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of GPCRs is an important mechanism for the regulation of some GPCRs. For example, in one form of signal transduction, the effect of hormone binding is the activation inside the cell of the enzyme, adenylate cyclase. Enzyme activation by hormones is dependent on the presence of the  
25 nucleotide GTP. GTP also influences hormone binding. A G protein connects the hormone receptor to adenylate cyclase. G protein exchanges GTP for bound GDP when activated by a hormone receptor. The GTP-carrying form then binds to activated adenylate cyclase. Hydrolysis of GTP to GDP, catalyzed by the G protein itself, returns the G protein to its basal, inactive form. Thus, the G protein serves a  
30 dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

- 3 -

Over the past 15 years, nearly 350 therapeutic agents targeting GPCRs have been successfully introduced onto the market. This indicates that these receptors have an established, proven history as therapeutic targets. Clearly, there is an on-going need for identification and characterization of further GPCRs which can play a role in preventing, ameliorating, or correcting dysfunctions or diseases including, but not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, pain, cancers, anorexia, bulimia, asthma, Parkinson's diseases, acute heart failure, hypotension, hypertension, urinary retention, osteoporosis, angina pectoris, myocardial infarction, ulcers, asthma, allergies, multiple sclerosis, benign prostatic hypertrophy, and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, several mental retardation, and dyskinesias, such as Huntington's disease and Tourett's syndrome.

15

#### P2Y Receptors

Extracellular nucleotides induce a wide variety of responses in many cell types, including muscle contraction and relaxation, vasodilation, neurotransmission, platelet aggregation, ion transport regulation, and cell growth. The effects are exerted mainly through two types of receptors: P2Y type G protein-coupled receptors, and P2X type ligand-gated ion channels. Nine distinct nucleotide-stimulated G protein-coupled receptors members have been characterized to date in humans. The nine receptors can be further subdivided into three groups according to ligand specificity: those activated by adenine nucleotides (P2Y1, P2Y11, P2Y12, and P2Y13), those activated by uridine nucleotides (P2Y4, P2Y6, CYSLT1, and GPR105), and those activated by both adenine and uridine nucleotides (P2Y2). The naturally occurring nucleotides that have been found to bind to these receptors have invariably been nucleotide diphosphates and nucleotide triphosphates. For example, ATP is the energy source for many biochemical reactions, a precursor for ribonucleic acid (RNA) synthesis, the precursor for cyclic AMP synthesis, etc. However, ATP also functions as an extracellular messenger in neuronal and non-neuronal tissues. Extracellular ATP exerts its

30

- 4 -

effects on these tissues by acting through membrane-associated purinoreceptors (Burnstock, G. Ann. NY Acad. Sci. (1990) 603:1-17) which can be either ligand-gated ion channels (Bean, B. P. (1992) Trends Pharmac. Sci. 12:87-90; Bean, B. P. and Fried, D. D. (1990) Ion Channels 2:169-203) that are generally referred to as P2X receptors, (but also known as: purinergic channels, P2X R-channels, and ATP-gated channels) or G-protein-coupled (P2Y) receptors (Barnard, E. A. et al. (1994) Trends Pharmac. Sci. 15:67-70). See U.S. Patent 5,856,129.

#### Adenosine receptors

In addition to nucleotides, nucleosides have also been shown to have extracellular signaling functions. Adenosine, a purine nucleoside, is a ubiquitous modulator of numerous physiological activities, particularly within the cardiovascular and nervous systems. The effects of adenosine appear to be mediated by specific cell surface receptor proteins. Adenosine modulates diverse physiological functions including induction of sedation, vasodilation, suppression of cardiac rate and contractility, inhibition of platelet aggregability, stimulation of gluconeogenesis and inhibition of lipolysis. In addition to its effects on adenylate cyclase, adenosine has been shown to open potassium channels, reduce flux through calcium channels, and inhibit or stimulate phosphoinositide turnover through receptor-mediated mechanisms (See for example, C. E. Muller and B. Stein "Adenosine Receptor Antagonists: Structures and Potential Therapeutic Applications," Current Pharmaceutical Design, 2:501 (1996) and C. E. Muller "A.sub.1-Adenosine Receptor Antagonists," Exp. Opin. Ther. Patents 7(5):419 (1997)).

Adenosine receptors belong to the superfamily of purine receptors which are currently subdivided into P.sub.1 (adenosine) and P.sub.2 (ATP, AMP OR ADENOSINE RECEPTOR LIGAND, and other nucleotides) receptors. Four receptor subtypes for the nucleoside adenosine have been cloned so far from various species including humans. Two receptor subtypes (A.sub.1 and A.sub.2a) exhibit affinity for adenosine in the nanomolar range while two other known subtypes A.sub.2b and A.sub.3 are low-affinity receptors, with affinity for adenosine in the low-micromolar

- 5 -

range. A.sub.1 and A.sub.3 adenosine receptor activation can lead to an inhibition of adenylate cyclase activity, while A.sub.2a and A.sub.2b activation causes a stimulation of adenylate cyclase.

5     Adenosine 5'-monophosphate (AMP) and Adenosine

All cells contain adenosine and adenine nucleotides, and many cell types have been shown to release adenosine or adenine nucleotides upon stimulation. For example, mast cells have been reported to release adenosine upon antigen challenge [Marquardt, D.L et al. (1984) Proc. Natl. Acad. Sci. U.S.A. 81:6192-6]. Activation of  
10     platelets [Jarvis, G.E. et al. (1996) Eur J Pharmacol 315:203-12], neutrophils [Madara, J.L. et al (1993) J. Clin. Invest. 91:2320-5]) and eosinophils [Resnick, M.B. et al. (1993)] has been reported to induce the release of AMP. Once released, adenine nucleotides also can be converted to adenosine by ectonucleotidase enzymes. Adenosine has been found to be increased in the bronchoalveolar lavage fluid  
15     (BALF) of inflamed airways and in particular is known to be present at higher concentrations in the BALF of patients with chronic inflammatory conditions of the lung, such as asthma and COPD (Driver, A.G. et al. (1993) Am. J. Respir. Dis. 148:91-7).

20     AMP or adenosine introduced artificially, either by inhalation or by instillation, into the respiratory tracts of patients with asthma and other inflammatory diseases of the lungs can cause immediate bronchoconstriction [Polosa, R. and Holgate, S.T. (1997) Thorax 52:919-23]. The response is significantly higher in patients with asthma than in other lung diseases, and is rarely seen in normal volunteers. Therefore it has  
25     recently been suggested that the response to AMP or adenosine can be used to differentiate asthma from other related diseases and that it can further be used as a specific marker of disease activity since responsiveness correlates well with inflammatory status. In light of the bronchoconstrictive activity of AMP and adenosine, it is clear that endogenously produced AMP and adenosine potentially  
30     play a significant role in the pathology of asthma and other inflammatory lung diseases. Indeed, two drugs that have bronchodilator effects in asthma, theophylline

- 6 -

and enprofylline, are thought to achieve their effects by blocking the adenosine-induced activation of inflammatory cells and the subsequent release of inflammatory mediators.

5     P2Y15

WO0214511 discloses human P2Y15 receptor, its amino acid sequence and nucleotide sequence as well as their regulation. The mouse ortholog of P2Y15 is in the accession number XP\_139267.

10    Asthma

Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the  
15    airways, 2) airway hyperresponsiveness caused by a decreased control of airway caliber, and 3) airway inflammation. Certain cells are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils, eosinophils, and other cells that bind IgE. These effector cells accumulate at the site of allergic reaction in the airways and  
20    release toxic products that contribute to the acute pathology and eventually to the tissue destruction related to the disorder. Other resident cells, such as smooth muscle cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may contribute to the pathology. While the airway obstruction of asthma, presenting clinically as an intermittent wheeze and  
25    shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually make asthma a chronic disabling disorder requiring long-term management.

30    Despite recent important advances in our understanding of the pathophysiology of asthma, the disease appears to be increasing in prevalence and severity (Gergen and

Weiss, *Am. Rev. Respir. Dis.* 146, 823-24, 1992). It is estimated that 30-40% of the population suffer with atopic allergy, and 15% of children and 5% of adults in the population suffer from asthma (Gergen and Weiss, 1992). Thus, an enormous burden is placed on our health care resources. However, both diagnosis and treatment of asthma are difficult. The severity of lung tissue inflammation is not easy to measure and the symptoms of the disease are often indistinguishable from those of respiratory infections, chronic respiratory inflammatory disorders, allergic rhinitis, or other respiratory disorders. Often, the inciting allergen cannot be determined, making removal of the causative environmental agent difficult. Current pharmacological treatments suffer their own set of disadvantages. Commonly used therapeutic agents, such as beta agonists, can act as symptom relievers to transiently improve pulmonary function, but do not affect the underlying inflammation. Agents that can reduce the underlying inflammation, such as anti-inflammatory steroids, can have major drawbacks that range from immunosuppression to bone loss (Goodman and Gilman's THE PHARMACOLOGIC BASIS OF THERAPEUTICS, Seventh Edition, MacMillan Publishing Company, NY, USA, 1985). In addition, many of the present therapies, such as inhaled corticosteroids, are short-lasting, inconvenient to use, and must be used often on a regular basis, in some cases for life, making failure of patients to comply with the treatment a major problem and thereby reducing their effectiveness as a treatment.

Because of the problems associated with conventional therapies, alternative treatment strategies have been evaluated. Glycophorin A (Chu and Sharom, *Cell. Immunol.* 145, 223-39, 1992), cyclosporin (Alexander *et al.*, *Lancet* 339, 324-28, 1992), and a nonapeptide fragment of IL-2 (Zav'yalov *et al.*, *Immunol. Lett.* 31, 285-88, 1992) all inhibit interleukin-2 dependent T lymphocyte proliferation; however, they are known to have many other effects. For example, cyclosporin is used as an immunosuppressant after organ transplantation. While these agents may represent alternatives to steroids in the treatment of asthmatics, they inhibit interleukin-2 dependent T lymphocyte proliferation and potentially critical immune functions associated with homeostasis. Other treatments that block the release or activity of



- 8 -

mediators of bronchoconstriction, such as cromones or anti-leukotrienes, have recently been introduced for the treatment of mild asthma, but they are expensive and not effective in all patients and it is unclear whether they have any effect on the chronic changes associated with asthmatic inflammation. There remains a need in the art for the identification of a treatment that can act in pathways critical to the development of asthma that blocks both the episodic attacks of the disorder and preferentially dampens the hyperactive allergic immune response without immunocompromising the patient.

#### 10 SUMMARY OF THE INVENTION

The present invention is based on the discovery that G protein coupled receptor GPR80 (Lee, DK et al., 2001, Gene 275:83-91) is regulated *in vivo* by AMP and adenosine. It is referred to as P2Y15 G protein-coupled receptor hereinafter.

15 It is an object of the invention to provide reagents and methods for regulating a human P2Y15 G protein-coupled receptor. This and other objects of the invention are provided by one or more of the embodiments described below.

20 One embodiment of the invention is a method for detecting the activity of P2Y15 in a sample, which method comprises the steps of: a) incubating a sample with P2Y15 and a ligand under conditions which allow binding of P2Y15 and the ligand, and b) detecting a second messenger, wherein said ligand is AMP or adenosine receptor ligand.

25 The method may further comprises the steps of: a) incubating a second sample with P2Y15 in the absence of the ligand under conditions which allow binding of P2Y15 and the ligand, and b) detecting a second messenger. The sample may comprise cells expressing P2Y15 or cell membranes bearing P2Y15.

30

Another embodiment of the invention is a method of screening for an agent that modulate P2Y15 activity using cells expressing P2Y15. The method comprises a) incubating a first sample of said cells in the presence of said agent and a second sample of said cells in the absence of said agent, both said samples under conditions  
5 which allow binding of AMP or adenosine receptor ligand to P2Y15; b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and c) comparing the results of said second messenger assays for said first and second samples.

10 Yet another embodiment of the invention is a method of screening for an agent to modulate P2Y15 activity using cell membranes bearing P2Y15. The method comprises a) incubating a first sample of said cell membranes in the presence of said agent and a second sample of said cell membranes in the absence of said agent, both  
15 said samples under conditions which allow binding of AMP or adenosine receptor ligand to P2Y15; b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and c) comparing the results of said second messenger assays for said first and second samples.

Further embodiment of the invention is a method for determining if a test compound  
20 increases or decreases the activity of P2Y15 using cells expressing P2Y15. The method comprises a) incubating a first sample of said cells in the presence of said test compound and a second sample of said cells in the absence of said test compound, both said samples under conditions which permit binding of AMP or adenosine receptor ligand to P2Y15; b) detecting a signalling activity of P2Y15  
25 polypeptide in said first and second samples, and c) comparing the results of said second messenger assays for said first and second samples.

Another embodiment of the invention is a method for determining if a test compound increases or decreases the activity of P2Y15 using cell membranes bearing P2Y15.  
30 The method comprises: a) incubating a first sample of said cell membranes in the presence of said test compound and a second sample of said cell membranes in the

absence of said test compound, both said samples under conditions which permit binding of AMP or adenosine receptor ligand to P2Y15; b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and c) comparing the results of said second messenger assays for said first and second samples.

5

Another embodiment of the invention is a method of identifying an agent that modulates the function of P2Y15. The method comprises: a) contacting a P2Y15 polypeptide in the presence and absence of an agent under conditions permitting the binding of said AMP or adenosine receptor ligand to said P2Y15 polypeptide; and b) measuring the binding of said P2Y15 polypeptide to said agent, relative to the binding in the absence of said agent. The agent which changes binding is identified as a potential therapeutic agent for decreasing or increasing the function of P2Y15. The measuring is performed using a method selected from label displacement, surface plasmon resonance, fluorescence resonance energy transfer, fluorescence quenching, and fluorescence polarization. The agent may be selected from the group consisting of a natural or synthetic peptide, a polypeptide, an antibody or antigen-binding fragment thereof, a lipid, a carbohydrate, a nucleic acid, and a small organic molecule. The step of measuring a signalling activity of the P2Y15 polypeptide comprises detecting a change in the level of a second messenger.

15  
20

The step of detecting a signalling activity may comprise measurement of guanine nucleotide binding or exchange, adenylate cyclase activity, cAMP, protein kinase C activity, phosphatidylinositol breakdown, diacylglycerol, inositol triphosphate, intracellular calcium, arachinoid acid concentration, MAP kinase activity, tyrosine kinase activity, and reporter gene expression.

25

Further embodiment of the invention is a reagent that modulates the activity of a P2Y15 polypeptide or polynucleotide. The reagent is identified by any of the above method.

30

Another embodiment of the invention is a pharmaceutical composition, which comprises the above mentioned reagent and a pharmaceutically acceptable carrier.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

5

Fig. 1 shows the relative expression of human P2Y15 GPCR in various human tissues and cells.

10

Fig. 2 shows calcium mobilization (A) and cyclic AMP generation (B) stimulated by AMP (filled symbols) and adenosine (empty symbols) in transfected HEK293 cells expressing P2Y15 (circles).

15

Fig. 3 shows specific binding of increasing concentrations of  $^3\text{H}$ -adenosine (A) and  $^{32}\text{P}$ -AMP (B) to transfected HEK293 cells expressing P2Y15 (circles) and nontransfected HEK293 cells (squares). The binding of  $^3\text{H}$ -AMP (C) or  $^3\text{H}$ -adenosine (D) to P2Y15-transfected cells and nontransfected cells could be competed in both cases by unlabeled AMP (filled symbols) or adenosine (open symbols).

20

Fig. 4 shows non-specific antagonists of adenosine block AMP- and adenosine-induced P2Y15 signaling. Calcium mobilization induced by  $10^{-6}$  M AMP (A) or  $10^{-6}$  M adenosine (B) was effectively blocked in a dose-dependent manner by theophylline, IBMX, 8-phenyltheophylline, caffeine, and AMP-CP.  $K_i$  values for each of the antagonists are shown.

25

Fig. 5 shows the relative intensity of expression of P2Y15 and various well known genes and markers expressed in mast cells as measured by DNA microarray analysis.

### **DETAILED DESCRIPTION OF THE INVENTION**

#### *Relevance of P2Y15 GPCR for the treatment of asthma*

30

P2Y15 GPCR is a seven-transmembrane-domain molecule that has highest homology to P2Y1. It was originally found in a search for P2Y homologs in genomic sequence databases. Only one EST has been reported to date for this gene, from a cDNA

library derived from normal human epithelium. The applicants performed expression profiling of this gene showing that it is expressed highest in the trachea, salivary glands, and kidneys, and less so in fetal brain, colon, placenta, and lung.

5        Although P2Y15 GPCR is closest in homology to P2Y1, which binds adenine nucleotides (ATP and AMP OR ADENOSINE RECEPTOR LIGAND), it also has significant homology to P2Y2 and P2Y4, which bind both A and U nucleotides, to P2Y3, which binds U nucleotides, and to leukotriene receptors, which bind LTB<sub>4</sub>, LTC<sub>4</sub>, and LTD<sub>4</sub>. The applicants therefore determined the true ligand of this  
10        receptor empirically.

In studies of airway epithelia, both ATP and UTP have been found to equipotently regulate epithelial electrolyte and water transport, trigger mucin secretion, and increase ciliary beat frequency. In the trachea, nucleotides can induce tracheal gland  
15        serous cells, which are responsible for the secretion of antibacterial and anti-proteolytic proteins, to produce secretory leukocyte proteinase inhibitor and to increase chloride transport. Studies in a mouse knockout of the P2Y2 receptor show that it is the dominant extracellular nucleotide receptor in airway epithelium, but that other nucleotide receptors exist that function similarly in the respiratory tract.

20        Expression profiling studies of P2Y15 GPCR performed by the applicants show that it is expressed highly in tissues of the upper respiratory tract. Its high expression in the salivary glands and trachea indicate that it plays a role in exocrine secretion, which in the airways has mainly a protective role. In asthma, however, over-  
25        production of mucin contributes to the viscid mucus plugs that occlude asthmatic airways. Submucosal glands in the large airways of asthmatics also frequently show evidence of hyperplasia, which may somehow be due to overstimulation by external mediators.

30        Both agonists and antagonists of the P2Y15 receptor can have a beneficial effect in asthmatics. Agonists can increase protective protein secretion and increase ciliary

beat rate, while antagonists can slow mucus production and glandular hyperplasia, prevent smooth muscle contraction, and reduce the sensitivity to respiratory tract irritants, including allergens, polluted, dry, or cold air, low oxygen concentrations, high CO<sub>2</sub> or CO concentrations, superoxides, and inflammatory mediators.

5

It has been discovered by the present applicant that the P2Y<sub>15</sub> GPCR can be regulated to treat bronchoconstriction and inflammation in diseases such as allergies including but not limited to asthma. Human P2Y<sub>15</sub> GPCR has the amino acid sequence shown in SEQ ID NO:2.

10

Human P2Y<sub>15</sub> GPCR also can be used to screen for human P2Y<sub>15</sub> GPCR agonists and antagonists.

*Relevance of P2Y<sub>15</sub> GPCR for the treatment of other diseases*

15

Within the kidney, analysis of P2Y<sub>15</sub> transcript localization by in situ hybridization performed by the applicants showed that expression appears to be highest in the kidney microvasculature. Since theophylline and caffeine, which act as inhibitors of P2Y<sub>15</sub> signaling, are known to have diuretic effects on the kidneys, other antagonists or agonists of P2Y<sub>15</sub> are also expected to have effects on kidney function. Such antagonists and agonists of P2Y<sub>15</sub> can be used to therapeutically regulate urine production, and in doing so, can also be beneficial in the regulation of the concentration of blood components, such as electrolytes and proteins. For example, P2Y<sub>15</sub> antagonists or agonists can be used in the treatment of diseases such as congestive heart failure, high blood pressure, edema, cirrhosis of the liver, and the nephrotic syndrome.

20  
25

In an analysis performed by the applicants using DNA microarrays to detect the expression levels of gene transcripts in human umbilical cord blood-derived mast cells, the expression of P2Y<sub>15</sub> transcripts was found to be, in terms of expression intensity, among the top 1% of genes expressed in these cells. Because of this extremely high expression, P2Y<sub>15</sub> is considered by the applicants to be an important

30

- 14 -

signaling molecule for mast cells. Consequently, regulation of P2Y<sub>15</sub> signaling can be used as a method to treat diseases and conditions in which mast cells are involved. Increased numbers of mast cells are found in many pathological conditions. For example, mast cell hyperplasia in the skin (mastocytosis) manifests with skin lesions and may present with symptoms of urticaria and flushing due to the chemical mediators released during mast cell degranulation. Children may develop single mastocytomas or the multiple cutaneous lesions of urticaria pigmentosa. In adults, multiple organ involvement can occur (notably affecting bone, liver, spleen and lymph nodes) even without apparent skin lesions (systemic mastocytosis). Lesions of the bone may be localised or widespread, osteoclastic or osteoblastic. Increased mast cell numbers are also seen in some inflammatory bowel diseases (ulcerative colitis, Crohn's disease) and in parasitic infections. Cutaneous neurofibromas, benign and malignant breast lesions, and some soft tissue tumours also show high numbers of mast cells. Mast cell numbers are also found to be high in interstitial cystitis and in idiopathic reduced bladder storage (sensory urge incontinence). Even in disease conditions where mast cell numbers are not recognized to be abnormally high, regulation of P2Y<sub>15</sub> signaling in mast cells can have a beneficial effect. For example, the degranulation of inflammatory cells has been reported to lead to the inactivation of muscarinic receptors on presynaptic nerve endings, interfering with the inhibitory function of the receptors and leading to an increased release of acetylcholine into the synapse. Such inactivation may lead to abnormal muscle contractions and instability, such as that seen in asthma and in overactive bladder. Preventing dysregulated degranulation of inflammatory cells, such as mast cells, by regulating P2Y<sub>15</sub> signaling can therefore be one way of treating symptomatic muscle contractions and instability.

Expression profiling studies of P2Y<sub>15</sub> GPCR performed by the applicants show that it is expressed highly in the salivary glands and trachea. This indicates that antagonists or agonists of P2Y<sub>15</sub> can be used to treat upper respiratory and oral diseases or conditions in which there is an abnormal production of mucus or saliva, such as Sjogren's syndrome, dry mouth, dental carries, post-nasal drip, and cough.

**Definition**

As used herein the term "P2Y15" refers to a polypeptide that is encoded by any  
5 polynucleotide selected from the group consisting of

- a) a polynucleotide encoding a P2Y15 polypeptide comprising an amino acid  
sequence selected from the group consisting of:
  - 10 amino acid sequences which are at least about 50% identical to the amino acid  
sequence shown in SEQ ID NO:2, 4, or 6; and  
the amino acid sequence shown in SEQ ID NO:2, 4, or 6
- b) a polynucleotide comprising the sequence of SEQ ID NO: 1, 3, or 5;
- 15 c) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and (b) and encodes a P2Y15 polypeptide;
- d) a polynucleotide the nucleic acid sequence of which deviates from the nucleic  
20 acid sequences specified in (a) to (c) due to the degeneration of the genetic  
code and encodes a P2Y15; and
- e) a polynucleotide, which represents a fragment, derivative or allelic variation  
of a nucleic acid sequence specified in (a) to (d) and encodes a P2Y15 and  
25 maintain P2Y15 activity.

As used herein the term "Ligand" refers to a molecule which binds to a receptor in a  
manner that is similar or equivalent to AMP.

30 As used herein the term "Adenosine receptor ligand" refers to specific and non-specific agonists and antagonists of adenosine receptors, including, but not limited to



- 16 -

adenosine, 2-chloroadenosine, N6-Cyclopentyladenosine (CPA), CGS-21680 hydrochloride, Chloro-IB-MECA, 5'-(N-Ethylcarboxamido) adenosine (NECA) and its derivatives, 8-Phenyltheophylline (8-PT), 3-isobutyl-1 methylxanthine (IBMX), Alloxazine, 8-(p-Sulfophenyl) theophylline (8-SPT), 8-cyclopentyl-1,3-dipropyl-  
5 xanthine (DPCPX), caffeine, theophylline, and enprofylline. Other adenosine receptor ligands are also known to those skilled in the art, e.g., as disclosed in Pharmacological reviews 53:527-552, 2001.

As used herein the term "Second messenger" refers to a molecule, generated or  
10 caused to vary in concentration by the activation of a G-Protein Coupled Receptor, that participates in the transduction of a signal from that GPCR. Non-limiting examples of second messengers include cAMP, diacylglycerol, inositol triphosphate, arachidonic acid metabolites, calcium ions. The term "change in the level of a second messenger" refers to an increase or decrease of at least 10% in the detected level of a  
15 given second messenger relative to the amount detected in an assay performed in the absence of a candidate modulator.

As used herein the term "Second messenger assay" preferably comprises the measurement of guanine nucleotide binding or exchange, adenylate cyclase, intra-  
20 cellular cAMP, intracellular inositol phosphate, intracellular diacylglycerol concentration, arachidonic acid concentration, calcium mobilization, MAP kinase(s) or tyrosine kinase(s), protein kinase C activity, or reporter gene expression or an aequorin-based assay according to methods known in the art and defined herein.

25 As used herein the term "Sample" is the source of molecules being tested for the presence of an agent or modulator compound that modulates binding to or signalling activity of a P2Y15 polypeptide. A sample can be an environmental sample, a natural extract of animal, plant yeast or bacterial cells or tissues, a clinical sample, a synthetic sample, or a conditioned medium from recombinant cells or a fermentation  
30 process.

- 17 -

As used herein, the term "membrane fraction" refers to a preparation of cellular lipid membranes comprising a P2Y<sub>15</sub> polypeptide. As the term is used herein, a "membrane fraction" is distinct from a cellular homogenate, in that at least a portion (i.e., at least 10%, and preferably more) of non-membrane-associated cellular constituents has been removed. The term "membrane associated" refers to those cellular constituents that are either integrated into a lipid membrane or are physically associated with a component that is integrated into a lipid membrane.

As used herein the term "Conditions which allow the binding of P2Y<sub>15</sub> and a ligand" refers to conditions of, for example, temperature, salt concentration, pH and protein concentration under which a ligand binds P2Y<sub>15</sub>. Exact binding conditions will vary depending upon the nature of the assay, for example, whether the assay uses viable cells or only membrane fraction of cells.

#### Polypeptides

P2Y<sub>15</sub> GPCR polypeptides according to the invention comprise at least 10, 12, 15, 20, 24, 30, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, or 350 contiguous amino acids selected from the amino acid sequence shown in SEQ ID NO:2 or a biologically active variant of that sequence, as defined below. A P2Y<sub>15</sub> GPCR polypeptide of the invention therefore can be a portion of a P2Y<sub>15</sub> GPCR, a full-length P2Y<sub>15</sub> GPCR, or a fusion protein comprising all or a portion of a P2Y<sub>15</sub> GPCR.

A phylogenetic analysis comparing the protein sequence with other G protein-coupled receptors (GPCRs) places the molecule among a cluster of other P2Y receptors, distant from the known receptors for adenosine (van den Berge, M., Kerstjens, H. A., and Postma, D. S. (2002) Clin Exp Allergy 32, 824-830). The gene sequence is found on the genomic contig NT\_009952 which has been localized to human chromosome 13q32.

30

Biologically Active Variants

P2Y<sub>15</sub> GPCR polypeptide variants which are biologically active, *i.e.*, retain the ability to bind a ligand to produce a biological effect, such as cyclic AMP formation, mobilization of intracellular calcium, or phosphoinositide metabolism, also are

5 P2Y<sub>15</sub> GPCR polypeptides. Preferably, naturally or non-naturally occurring P2Y<sub>15</sub> GPCR polypeptide variants have amino acid sequences which are at least about 50, 55, 60, 65, 70, more preferably about 75, 90, 96, or 98% identical to an amino acid sequence shown in SEQ ID NO:2 or a fragment thereof. Percent identity between a putative P2Y<sub>15</sub> GPCR polypeptide variant and an amino acid sequence of SEQ ID

10 NO:2 is determined by conventional methods. See, for example, Altschul *et al.*, *Bull. Math. Bio.* 48:603 (1986), and Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1992). Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff & Henikoff, 1992.

15

Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson & Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino

20 acid sequence of a putative variant. The FASTA algorithm is described by Pearson & Lipman, *Proc. Nat'l Acad. Sci. USA* 85:2444(1988), and by Pearson, *Meth. Enzymol.* 183:63 (1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (*e.g.*, SEQ ID NO: 2) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or

25 pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several

30 regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence the ktup value), then the trimmed

- 19 -

initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman & Wunsch, *J. Mol. Biol.* 48:444 (1970); Sellers, *SIAM J. Appl. Math.* 26:787 (1974)), which allows for amino acid insertions and deletions. Preferred parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, *Meth. Enzymol.* 183:63 (1990).

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as default.

Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity of a P2Y<sub>15</sub> GPCR polypeptide can be found using computer programs well known in the art, such as DNASTAR software. Whether an amino acid change results in a biologically active P2Y<sub>15</sub> GPCR polypeptide can readily be determined by assaying for binding to a ligand or by

- 20 -

conducting a functional assay, as described for example, in the specific Examples, below.

### Fusion Proteins

5 Fusion proteins are useful for generating antibodies against P2Y<sub>15</sub> GPCR polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of a P2Y<sub>15</sub> GPCR polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display  
10 systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

A P2Y<sub>15</sub> GPCR polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises  
15 at least 10, 12, 15, 20, 24, 30, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, or 325 contiguous amino acids of SEQ ID NO:2 or a biologically active variant of SEQ ID NO:2. Contiguous amino acids for use in a fusion protein can be selected from the amino acid sequence shown in SEQ ID NO:2 or from a biologically active variant of those sequences, such as those described above. The first polypeptide  
20 segment also can comprise full-length P2Y<sub>15</sub> G protein-coupled receptor.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include  $\beta$ -galactosidase,  $\beta$ -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including  
25 blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose  
30 binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions.

A fusion protein also can be engineered to contain a cleavage site located between the P2Y15 GPCR polypeptide-encoding sequence and the heterologous protein sequence, so that the P2Y15 GPCR polypeptide can be cleaved and purified away from the heterologous moiety.

5

A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences of SEQ ID NO:1 in proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Bio-technology (Santa Cruz, CA), MBL International Corporation (Woburn, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

10

15

#### Identification of Species Homologs

Species homologs of human P2Y15 GPCR polypeptide can be obtained using P2Y15 GPCR polynucleotides (described below) to make suitable probes or primers for screening cDNA expression libraries from other species, such as guinea pigs, monkeys, or yeast, identifying cDNAs which encode homologs of P2Y15 GPCR polypeptide, and expressing the cDNAs as is known in the art.

20

25

Species homologs of human P2Y15 GPCR polypeptide can also be obtained by performing a search, using the human or other species's P2Y15 polypeptide sequences as a query, in the Genbank database of the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov>) using the program tblastn. Mouse ortholog of P2Y15 can be found under accession number XP\_139267.

30

### Polynucleotides

A P2Y15 GPCR polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a P2Y15 GPCR polypeptide. A nucleotide sequence encoding SEQ ID NO:2 is shown in SEQ ID NO:1.

5

Degenerate nucleotide sequences encoding human P2Y15 GPCR polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, or 70, more preferably about 75, 90, 96, or 98% identical to a nucleotide sequence shown in SEQ ID NOS:1, 3, or 5 or its complement also are P2Y15 GPCR polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologs, and variants of P2Y15 GPCR polynucleotides which encode biologically active P2Y15 GPCR polypeptides also are P2Y15 GPCR polynucleotides.

10

15

### Identification of Polynucleotide Variants and Homologs

Variants and homologs of the P2Y15 GPCR polynucleotides described above also are P2Y15 GPCR polynucleotides. Typically, homologous P2Y15 GPCR polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known P2Y15 GPCR polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions--2X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2X SSC, 0.1% SDS, 50°C once, 30 minutes; then 2X SSC, room temperature twice, 10 minutes each--homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous nucleic acid strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

20

25

30

Species homologs of the P2Y15 GPCR polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression

libraries from other species, such as mice, monkeys, or yeast. Human variants of P2Y<sub>15</sub> GPCR polynucleotides can be identified, for example, by screening human cDNA expression libraries. It is well known that the  $T_m$  of a double-stranded DNA decreases by 1-1.5°C with every 1% decrease in homology (Bonner *et al.*, *J. Mol. Biol.* 81, 123 (1973). Variants of human P2Y<sub>15</sub> GPCR polynucleotides or P2Y<sub>15</sub> GPCR polynucleotides of other species can therefore be identified by hybridizing a putative homologous P2Y<sub>15</sub> GPCR polynucleotide with a polynucleotide having a nucleotide sequence of SEQ ID NO:1, 3 or 5 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of basepair mismatches within the test hybrid is calculated.

Nucleotide sequences which hybridize to P2Y<sub>15</sub> GPCR polynucleotides or their complements following stringent hybridization and/or wash conditions also are P2Y<sub>15</sub> GPCR polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989, at pages 9.50-9.51.

Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20 °C below the calculated  $T_m$  of the hybrid under study. The  $T_m$  of a hybrid between a P2Y<sub>15</sub> GPCR polynucleotide having a nucleotide sequence shown in SEQ ID NO:1, 3, or 5 or the complement thereof and a polynucleotide sequence which is at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation of Bolton and McCarthy, *Proc. Natl. Acad. Sci. U.S.A.* 48, 1390 (1962):

$$T_m = 81.5^\circ\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/l,$$

where  $l$  = the length of the hybrid in basepairs.



Stringent wash conditions include, for example, 4X SSC at 65°C, or 50% formamide, 4X SSC at 42°C, or 0.5X SSC, 0.1% SDS at 65°C. Highly stringent wash conditions include, for example, 0.2X SSC at 65 °C.

5     Preparation of Polynucleotides

A P2Y15 GPCR polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by  
10     using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated P2Y15 GPCR polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprises P2Y15 GPCR nucleotide sequences. Isolated polynucleotides are in preparations  
15     which are free or at least 70, 80, or 90% free of other molecules.

P2Y15 GPCR cDNA molecules can be made with standard molecular biology techniques, using P2Y15 GPCR mRNA as a template. P2Y15 GPCR cDNA molecules can thereafter be replicated using molecular biology techniques known in the  
20     art and disclosed in manuals such as Sambrook *et al.* (1989). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize P2Y15  
25     GPCR polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a P2Y15 GPCR polypeptide having, for example, the amino acid sequence shown in SEQ ID NO:2, 4, or 6 or a biologically active variant thereof.

Extending Polynucleotides

Various PCR-based methods can be used to extend the nucleic acid sequences encoding the disclosed portions of human P2Y<sub>15</sub> GPCR polypeptide to detect upstream sequences such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, *PCR Methods Applic.* 2, 318-322, 1993). Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region (Triglia *et al.*, *Nucleic Acids Res.* 16, 8186, 1988). Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72 °C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom *et al.*, *PCR Methods Applic.* 1, 111-119, 1991). In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

Another method which can be used to retrieve unknown sequences is that of Parker *et al.*, *Nucleic Acids Res.* 19, 3055-3060, 1991). Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif.) can be used to

walk genomic DNA (CLONTECH, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

5 When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

10

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser  
15 activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable  
20 for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

#### Obtaining Polypeptides

P2Y<sub>15</sub> GPCR polypeptides can be obtained, for example, by purification from cells,  
25 by expression of P2Y<sub>15</sub> GPCR polynucleotides, or by direct chemical synthesis.

#### Protein Purification

P2Y<sub>15</sub> GPCR polypeptides can be purified from any cell which expresses the receptor, including host cells which have been transfected with P2Y<sub>15</sub> GPCR  
30 polynucleotides which express such polypeptides. A purified P2Y<sub>15</sub> GPCR polypeptide is separated from other compounds which normally associate with the P2Y<sub>15</sub>

GPCR polypeptide in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

5

A P2Y<sub>15</sub> GPCR polypeptide can be conveniently isolated as a complex with its associated G protein, as described in the specific examples, below. A preparation of purified P2Y<sub>15</sub> GPCR polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

10

#### Expression of Polynucleotides

To express a P2Y<sub>15</sub> GPCR polypeptide, a P2Y<sub>15</sub> GPCR polynucleotide can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding P2Y<sub>15</sub> GPCR polypeptides and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described, for example, in Sambrook *et al.* (1989) and in Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1989.

15

20

A variety of expression vector/host systems can be utilized to contain and express sequences encoding a P2Y<sub>15</sub> GPCR polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

30

- 28 -

The control elements or regulatory sequences are those non-translated regions of the vector -- enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a P2Y<sub>15</sub> GPCR polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

#### Bacterial and Yeast Expression Systems

In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the P2Y<sub>15</sub> GPCR polypeptide. For example, when a large quantity of a P2Y<sub>15</sub> GPCR polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the P2Y<sub>15</sub> GPCR polypeptide can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced. pIN vectors (Van Heeke & Schuster, *J. Biol. Chem.* 264, 5503-5509, 1989) or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion

- 29 -

proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel *et al.* (1989) and Grant *et al.*, *Methods Enzymol.* 153, 516-544, 1987.

#### Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding P2Y15 GPCR polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6, 307-311, 1987). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used (Coruzzi *et al.*, *EMBO J.* 3, 1671-1680, 1984; Broglie *et al.*, *Science* 224, 838-843, 1984; Winter *et al.*, *Results Probl. Cell Differ.* 17, 85-105, 1991). These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (e.g., Hobbs or Murray, in MCGRAW HILL YEARBOOK OF SCIENCE AND TECHNOLOGY, McGraw Hill, New York, N.Y., pp. 191-196, 1992).

An insect system also can be used to express a P2Y15 GPCR polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding P2Y15 GPCR polypeptides can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of P2Y15

- 30 -

GPCR polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which P2Y15 GPCR polypeptides can be expressed (Engelhard *et al.*, *Proc. Nat. Acad. Sci.* 91, 3224-3227, 1994).

5

#### Mammalian Expression Systems

A number of viral-based expression systems can be used to express P2Y15 GPCR polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding P2Y15 GPCR polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing a P2Y15 GPCR polypeptide in infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci.* 81, 3655-3659, 1984). If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (*e.g.*, liposomes, polycationic amino polymers, or vesicles).

Specific initiation signals also can be used to achieve more efficient translation of sequences encoding P2Y15 GPCR polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a P2Y15 GPCR polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous

translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used (see Scharf *et al.*, *Results Probl. Cell Differ.* 20, 125-162, 1994).

5

#### Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed P2Y15 GPCR polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (*e.g.*, CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

10  
15

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express P2Y15 GPCR polypeptides can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced P2Y15 GPCR sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. See, for example, ANIMAL CELL CULTURE, R.I. Freshney, ed., 1986.

20  
25

30



- 32 -

Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, *Cell* 11, 223-32, 1977) and adenine phosphoribosyltransferase (Lowy *et al.*, *Cell* 22, 817-23, 1980) genes which can be employed in *tk* or *aprt* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate (Wigler *et al.*, *Proc. Natl. Acad. Sci.* 77, 3567-70, 1980), *npt* confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin *et al.*, *J. Mol. Biol.* 150, 1-14, 1981), and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murray, 1992, *supra*). Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci.* 85, 8047-51, 1988). Visible markers such as anthocyanins,  $\beta$ -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes *et al.*, *Methods Mol. Biol.* 55, 121-131, 1995).

#### Detecting Expression of Polypeptides

Although the presence of marker gene expression suggests that the P2Y15 GPCR polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding a P2Y15 GPCR polypeptide is inserted within a marker gene sequence, transformed cells containing sequences which encode a P2Y15 GPCR polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a P2Y15 GPCR polypeptide under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the P2Y15 GPCR polynucleotide.

Alternatively, host cells which contain a P2Y15 GPCR polynucleotide and which express a P2Y15 GPCR polypeptide can be identified by a variety of procedures

known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding a P2Y<sub>15</sub> GPCR polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a P2Y<sub>15</sub> GPCR polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding a P2Y<sub>15</sub> GPCR polypeptide to detect transformants which contain a P2Y<sub>15</sub> GPCR polynucleotide.

A variety of protocols for detecting and measuring the expression of a P2Y<sub>15</sub> GPCR polypeptide, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on a P2Y<sub>15</sub> GPCR polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton *et al.*, SEROLOGICAL METHODS: A LABORATORY MANUAL, APS Press, St. Paul, Minn., 1990) and Maddox *et al.*, *J. Exp. Med.* 158, 1211-1216, 1983).

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding P2Y<sub>15</sub> GPCR polypeptides include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding a P2Y<sub>15</sub> GPCR polypeptide can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3,

- 34 -

or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

#### Expression and Purification of Polypeptides

Host cells transformed with nucleotide sequences encoding a P2Y<sub>15</sub> GPCR polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode P2Y<sub>15</sub> GPCR polypeptides can be designed to contain signal sequences which direct secretion of soluble P2Y<sub>15</sub> GPCR polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound P2Y<sub>15</sub> GPCR polypeptide.

As discussed above, other constructions can be used to join a sequence encoding a P2Y<sub>15</sub> GPCR polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the P2Y<sub>15</sub> GPCR polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a P2Y<sub>15</sub> GPCR polypeptide and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography, as described in Porath *et al.*, *Prot.*

*Exp. Purif.* 3, 263-281, 1992), while the enterokinase cleavage site provides a means for purifying the P2Y<sub>15</sub> GPCR polypeptide from the fusion protein. Vectors which contain fusion proteins are disclosed in Kroll *et al.*, *DNA Cell Biol.* 12, 441-453, 1993.

5

#### Chemical Synthesis

Sequences encoding a P2Y<sub>15</sub> GPCR polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers *et al.*, *Nucl. Acids Res. Symp. Ser.* 215-223, 1980; Horn *et al.* *Nucl. Acids Res. Symp. Ser.* 225-232, 1980). Alternatively, a P2Y<sub>15</sub> GPCR polypeptide itself can be produced using  
10 chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques (Merrifield, *J. Am. Chem. Soc.* 85, 2149-2154, 1963; Roberge *et al.*, *Science* 269, 202-204, 1995). Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be  
15 achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of P2Y<sub>15</sub> GPCR polypeptides can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high  
20 performance liquid chromatography (*e.g.*, Creighton, *PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES*, WH Freeman and Co., New York, N.Y., 1983). The composition of a synthetic P2Y<sub>15</sub> GPCR polypeptide can be confirmed by amino acid analysis or sequencing (*e.g.*, the Edman degradation procedure; *see* Creighton, *supra*). Additionally, any portion of the amino acid sequence of the P2Y<sub>15</sub> GPCR  
25 polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

#### Production of Altered Polypeptides

30 As will be understood by those of skill in the art, it may be advantageous to produce P2Y<sub>15</sub> GPCR polypeptide-encoding nucleotide sequences possessing non-naturally

occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

5

The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter P2Y<sub>15</sub> GPCR polypeptide-encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

15

### Orthologs

Orthologs, such as the mouse ortholog of P2Y<sub>15</sub> GPCR (GenBank accession number XP\_139267.1; SEQ ID NO: 3/amino acid seq; SEQ ID NO:4 ), may also be produced. The mouse ortholog has an amino acid sequence that is 85% identical and 88% homologous to the human P2Y<sub>15</sub> GPCR amino acid sequence. It will be appreciated by one of skill in the art that due to the high degree of identity between the mouse and the human proteins that the mouse can be used as a model system to screen for and study the effect of target compounds that effect P2Y<sub>15</sub> GPCR activity. The mouse can also be used as a model system to study the effect of mutations in the P2Y<sub>15</sub> gene or over- or under-expression of P2Y<sub>15</sub> GPCR on disease, including but not limited to allergic responses. The mouse ortholog of P2Y<sub>15</sub> GPCR is also useful as a purified protein to screen for target compounds that modulate receptor activity. This protein may also be purified to aid in efforts to solve the structure of P2Y<sub>15</sub> GPCR and the design of small molecules that effect protein activity. The rat ortholog of P2Y<sub>15</sub> was found by applicant by using the mouse protein sequence in a tblastn query against the rat genome trace sequences subset of Genbank. Seven

30

- 37 -

transmembrane regions as predicted by the computer program TMpred are indicated with heavy overlines and numbered TM1-7. The Genbank accession number of the rat P2Y15 sequence is AY191367 (SEQ ID NO:5/amino acid seq;SEQ ID NO:6).

5     Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of a P2Y15 GPCR polypeptide. "Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv, which are capable of binding an epitope of a P2Y15 GPCR polypeptide.  
10     Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acids.

An antibody which specifically binds to an epitope of a P2Y15 GPCR polypeptide  
15     can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immuno-precipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in  
20     the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the immunogen.

Typically, an antibody which specifically binds to a P2Y15 GPCR polypeptide  
25     provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to P2Y15 GPCR polypeptides do not detect other proteins in immunochemical assays and can immunoprecipitate a P2Y15 GPCR polypeptide from solution.

30

- 38 -

P2Y<sub>15</sub> GPCR polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a P2Y<sub>15</sub> GPCR polypeptide can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies which specifically bind to a P2Y<sub>15</sub> GPCR polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler *et al.*, *Nature* 256, 495-497, 1985; Kozbor *et al.*, *J. Immunol. Methods* 81, 31-42, 1985; Cote *et al.*, *Proc. Natl. Acad. Sci.* 80, 2026-2030, 1983; Cole *et al.*, *Mol. Cell Biol.* 62, 109-120, 1984).

In addition, techniques developed for the production of "chimeric antibodies," the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (Morrison *et al.*, *Proc. Natl. Acad. Sci.* 81, 6851-6855, 1984; Neuberger *et al.*, *Nature* 312, 604-608, 1984; Takeda *et al.*, *Nature* 314, 452-454, 1985). Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Alternatively,

- 39 -

humanized antibodies can be produced using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to a P2Y<sub>15</sub> GPCR polypeptide can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

5

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to P2Y<sub>15</sub> GPCR polypeptides. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton, *Proc. Natl. Acad. Sci.* 88, 11120-23, 1991).

10

Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template (Thirion *et al.*, 1996, *Eur. J. Cancer Prev.* 5, 507-11). Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma & Morrison, 1997, *Nat. Biotechnol.* 15, 159-63. Construction of bivalent, bispecific single-chain antibodies is taught in Mallender & Voss, 1994, *J. Biol. Chem.* 269, 199-206.

15

20

A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology (Verhaar *et al.*, 1995, *Int. J. Cancer* 61, 497-501; Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91).

25

Antibodies which specifically bind to P2Y<sub>15</sub> GPCR polypeptides also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as

30



disclosed in the literature (Orlandi *et al.*, *Proc. Natl. Acad. Sci.* 86, 3833-3837, 1989; Winter *et al.*, *Nature* 349, 293-299, 1991).

5 Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

10 Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a P2Y<sub>15</sub> GPCR polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

15 Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide  
20 is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of P2Y<sub>15</sub> GPCR gene products in the cell.

25 Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such  
30 alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate,

carboxymethyl esters, carbonates, and phosphate triesters. See Brown, *Meth. Mol. Biol.* 20, 1-8, 1994; Sonveaux, *Meth. Mol. Biol.* 26, 1-72, 1994; Uhlmann *et al.*, *Chem. Rev.* 90, 543-583, 1990.

5 Modifications of P2Y15 GPCR gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the P2Y15 GPCR. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology.  
10 Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature (*e.g.*, Gee *et al.*, in Huber & Carr, MOLECULAR AND IMMUNOLOGIC APPROACHES, Futura Publishing Co., Mt. Kisco, N.Y., 1994). An antisense  
15 oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a P2Y15 GPCR  
20 polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a P2Y15 GPCR polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent P2Y15 GPCR nucleotides, can provide sufficient targeting specificity for P2Y15 GPCR mRNA. Preferably, each stretch of  
25 complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular P2Y15  
30 GPCR polynucleotide sequence.

Antisense oligonucleotides can be modified without affecting their ability to hybridize to a P2Y<sub>15</sub> GPCR polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art. See, e.g., Agrawal *et al.*, *Trends Biotechnol.* 10, 152-158, 1992; Uhlmann *et al.*, *Chem. Rev.* 90, 543-584, 1990; Uhlmann *et al.*, *Tetrahedron. Lett.* 215, 3539-3542, 1987.

#### Ribozymes

Ribozymes are RNA molecules with catalytic activity. See, e.g., Cech, *Science* 236, 1532-1539; 1987; Cech, *Ann. Rev. Biochem.* 59, 543-568; 1990, Cech, *Curr. Opin. Struct. Biol.* 2, 605-609; 1992, Couture & Stinchcomb, *Trends Genet.* 12, 510-515, 1996. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff *et al.*, U.S. Patent 5,641,673). The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences.

The coding sequence of a P2Y<sub>15</sub> GPCR polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from the P2Y<sub>15</sub> GPCR polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art (see Haseloff *et al.* *Nature* 334, 585-591, 1988). For example, the cleavage activity of ribozymes can be targeted to specific RNAs by

engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target (see, for example, Gerlach *et al.*, EP 321,201).

5 Specific ribozyme cleavage sites within a P2Y<sub>15</sub> GPCR RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which  
10 may render the target inoperable. Suitability of candidate P2Y<sub>15</sub> GPCR RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally  
15 related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or  
20 calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease P2Y<sub>15</sub> GPCR expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding  
25 DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

As taught in Haseloff *et al.*, U.S. Patent 5,641,673, ribozymes can be engineered so  
30 that ribozyme expression will occur in response to factors which induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of

regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

#### Differentially Expressed Genes

5 Described herein are methods for the identification of genes whose products interact with human P2Y<sub>15</sub> G protein-coupled receptor. Such genes may represent genes that are differentially expressed in disorders including, but not limited to, CNS disorders, cardiovascular disorders, asthma, osteoporosis, diabetes, and COPD. Further, such genes may represent genes that are differentially regulated in response to mani-  
10 pulations relevant to the progression or treatment of such diseases. Additionally, such genes may have a temporally modulated expression, increased or decreased at different stages of tissue or organism development. A differentially expressed gene may also have its expression modulated under control versus experimental conditions. In addition, the human P2Y<sub>15</sub> G protein-coupled receptor gene or gene  
15 product may itself be tested for differential expression.

The degree to which expression differs in a normal versus a diseased state need only be large enough to be visualized via standard characterization techniques such as differential display techniques. Other such standard characterization techniques by  
20 which expression differences may be visualized include but are not limited to, quantitative RT (reverse transcriptase), PCR, and Northern analysis.

#### Identification of Differentially Expressed Genes

To identify differentially expressed genes total RNA or, preferably, mRNA is  
25 isolated from tissues of interest. For example, RNA samples are obtained from tissues of experimental subjects and from corresponding tissues of control subjects. Any RNA isolation technique that does not select against the isolation of mRNA may be utilized for the purification of such RNA samples. See, for example, Ausubel *et al.*, ed., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, Inc.  
30 New York, 1987-1993. Large numbers of tissue samples may readily be processed

using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, U.S. Patent 4,843,155.

5 Transcripts within the collected RNA samples that represent RNA produced by differentially expressed genes are identified by methods well known to those of skill in the art. They include, for example, differential screening (Tedder *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 85, 208-12, 1988), subtractive hybridization (Hedrick *et al.*, *Nature* 308, 149-53; Lee *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 88, 2825, 1984), and differential display (Liang & Pardee, *Science* 257, 967-71, 1992; U.S. Patent 10 5,262,311), and microarrays.

The differential expression information may itself suggest relevant methods for the treatment of disorders involving the human P2Y15 G protein-coupled receptor. For example, treatment may include a modulation of expression of the differentially 15 expressed genes and/or the gene encoding the human P2Y15 G protein-coupled receptor. The differential expression information may indicate whether the expression or activity of the differentially expressed gene or gene product or the human P2Y15 G protein-coupled receptor gene or gene product are up-regulated or down-regulated.

20

#### Screening Methods

The invention provides assays for screening test compounds which bind to or modulate the activity of a P2Y15 GPCR polypeptide or a P2Y15 GPCR polynucleotide. A test compound preferably binds to a P2Y15 GPCR polypeptide or 25 polynucleotide. More preferably, a test compound decreases or increases a biological effect mediated via human P2Y15 GPCR by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the test compound.

#### Test Compounds

30 Test compounds can be pharmacologic agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The com-

pounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinantly, or synthesized by chemical methods known in the art. If desired, test compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially addressable parallel solid phase or solution phase libraries, synthetic library methods requiring deconvolution, the "one-bead one-compound" library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds. See Lam, *Anticancer Drug Des.* 12, 145, 1997.

Methods for the synthesis of molecular libraries are well known in the art (see, for example, DeWitt *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 90, 6909, 1993; Erb *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 91, 11422, 1994; Zuckermann *et al.*, *J. Med. Chem.* 37, 2678, 1994; Cho *et al.*, *Science* 261, 1303, 1993; Carell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33, 2059, 1994; Carell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33, 2061; Gallop *et al.*, *J. Med. Chem.* 37, 1233, 1994). Libraries of compounds can be presented in solution (see, e.g., Houghten, *BioTechniques* 13, 412-421, 1992), or on beads (Lam, *Nature* 354, 82-84, 1991), chips (Fodor, *Nature* 364, 555-556, 1993), bacteria or spores (Ladner, U.S. Patent 5,223,409), plasmids (Cull *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 89, 1865-1869, 1992), or phage (Scott & Smith, *Science* 249, 386-390, 1990; Devlin, *Science* 249, 404-406, 1990); Cwirla *et al.*, *Proc. Natl. Acad. Sci.* 97, 6378-6382, 1990; Felici, *J. Mol. Biol.* 222, 301-310, 1991; and Ladner, U.S. Patent 5,223,409).

#### High Throughput Screening

Test compounds can be screened for the ability to bind to P2Y<sub>15</sub> GPCR polypeptides or polynucleotides or to affect P2Y<sub>15</sub> GPCR activity or P2Y<sub>15</sub> GPCR gene expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well

microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 50 to 500  $\mu$ l. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the 96-well format.

5

Alternatively, "free format assays," or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by Jayawickreme *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 19, 1614-18 (1994). The cells are placed under agarose in petri dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualized as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

10

15

Another example of a free format assay is described by Chelsky, "Strategies for Screening Combinatorial Libraries: Novel and Traditional Approaches," reported at the First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa. (Nov. 7-10, 1995). Chelsky placed a simple homogenous enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel. Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV-light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

20

25

Yet another example is described by Salmon *et al.*, *Molecular Diversity* 2, 57-63 (1996). In this example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar.

30

Another high throughput screening method is described in Beutel *et al.*, U.S. Patent 5,976,813. In this method, test samples are placed in a porous matrix. One or more



assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

5

#### Binding Assays

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of the P2Y<sub>15</sub> GPCR polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. In addition to adenosine and adenosine monophosphate, other potential ligands which may bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known GPCRs and analogues or derivatives thereof. Natural ligands of GPCRs include adrenomedullin, amylin, calcitonin gene related protein (CGRP), calcitonin, anandamide, serotonin, histamine, adrenalin, noradrenalin, platelet activating factor, thrombin, C5a, bradykinin, and chemokines. Other potential ligands which may bind to a polypeptide of the invention include, but are not limited to, specific and non-specific agonists and antagonists of Adenosine receptors, including caffeine, theophylline, enprofylline, and IBMX.

20

In binding assays, either the test compound or the P2Y<sub>15</sub> GPCR polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to the P2Y<sub>15</sub> GPCR polypeptide can then be accomplished, for example, by direct counting of radio-emission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

25

Alternatively, binding of a test compound to a P2Y<sub>15</sub> GPCR polypeptide can be determined without labeling either of the interactants. For example, a micro-

30

physiometer can be used to detect binding of a test compound with a P2Y<sub>15</sub> GPCR polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate  
5 can be used as an indicator of the interaction between a test compound and a P2Y<sub>15</sub> GPCR polypeptide (McConnell *et al.*, *Science* 257, 1906-1912, 1992).

Determining the ability of a test compound to bind to a P2Y<sub>15</sub> GPCR polypeptide also can be accomplished using a technology such as real-time Bimolecular  
10 Interaction Analysis (BIA) (Sjolander & Urbaniczky, *Anal. Chem.* 63, 2338-2345, 1991, and Szabo *et al.*, *Curr. Opin. Struct. Biol.* 5, 699-705, 1995). BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between  
15 biological molecules.

In yet another aspect of the invention, a P2Y<sub>15</sub> GPCR polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent 5,283,317; Zervos *et al.*, *Cell* 72, 223-232, 1993; Madura *et al.*, *J. Biol. Chem.* 268,  
20 12046-12054, 1993; Bartel *et al.*, *BioTechniques* 14, 920-924, 1993; Iwabuchi *et al.*, *Oncogene* 8, 1693-1696, 1993; and Brent W094/10300), to identify other proteins which bind to or interact with the P2Y<sub>15</sub> GPCR polypeptide and modulate its activity.

25 The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a P2Y<sub>15</sub> GPCR polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other  
30 construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known

transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with the P2Y15 GPCR polypeptide.

It may be desirable to immobilize either the P2Y15 GPCR polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the P2Y15 GPCR polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach the P2Y15 GPCR polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a P2Y15 GPCR polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, the P2Y15 GPCR polypeptide is a fusion protein comprising a domain that allows the P2Y15 GPCR polypeptide to be bound to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the

non-adsorbed P2Y15 GPCR polypeptide; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined  
5 either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a P2Y15  
10 GPCR polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated P2Y15 GPCR polypeptides (or polynucleotides) or test compounds can be prepared from biotin-NHS(N-hydroxysuccinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of  
15 streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a P2Y15 GPCR polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of the P2Y15 GPCR polypeptide, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

20 Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to the P2Y15 GPCR polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of the P2Y15 GPCR  
25 polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a P2Y15 GPCR polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a P2Y15 GPCR polypeptide or polynucleotide can be used in a cell-based assay  
30 system. A P2Y15 GPCR polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test

compound to a P2Y<sub>15</sub> GPCR polypeptide or polynucleotide is determined as described above.

Functional Assays

5 Test compounds can be tested for the ability to increase or decrease a biological effect of a P2Y<sub>15</sub> GPCR polypeptide. Such biological effects can be determined using the functional assays described in the specific examples, below. Functional assays can be carried out after contacting either a purified P2Y<sub>15</sub> GPCR polypeptide, a cell membrane preparation, or an intact cell with a test compound. A test com-  
10 pound which decreases a functional activity of a P2Y<sub>15</sub> GPCR by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing P2Y<sub>15</sub> GPCR activity. A test compound which increases P2Y<sub>15</sub> GPCR activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing  
15 P2Y<sub>15</sub> GPCR activity.

One such screening procedure involves the use of melanophores which are transfected to express a P2Y<sub>15</sub> GPCR polypeptide. Such a screening technique is described in WO 92/01810 published Feb. 6, 1992. Thus, for example, such an assay  
20 may be employed for screening for a compound which inhibits activation of the receptor polypeptide by contacting the melanophore cells which comprise the receptor with both a receptor ligand and a test compound to be screened. Inhibition of the signal generated by the ligand indicates that a test compound is a potential antagonist for the receptor, *i.e.*, inhibits activation of the receptor. The screen may be  
25 employed for identifying a test compound which activates the receptor by contacting such cells with compounds to be screened and determining whether each test compound generates a signal, *i.e.*, activates the receptor.

Other screening techniques include the use of cells which express a human P2Y<sub>15</sub> GPCR polypeptide (for example, transfected CHO cells) in a system which measures  
30 extracellular pH changes caused by receptor activation (*see, e.g., Science* 246,

- 53 -

181-296, 1989). For example, test compounds may be contacted with a cell which expresses a human P2Y<sub>15</sub> GPCR polypeptide and a second messenger response, *e.g.*, signal transduction or pH changes, can be measured to determine whether the test compound activates or inhibits the receptor.

5

Another such screening technique involves introducing RNA encoding a human P2Y<sub>15</sub> GPCR polypeptide into *Xenopus* oocytes to transiently express the receptor. The transfected oocytes can then be contacted with the receptor ligand and a test compound to be screened, followed by detection of inhibition or activation of a calcium signal in the case of screening for test compounds which are thought to inhibit activation of the receptor.

10

15

Another screening technique involves expressing a human P2Y<sub>15</sub> GPCR polypeptide in cells in which the receptor is linked to a phospholipase C or D. Such cells include endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening may be accomplished as described above by quantifying the degree of activation of the receptor from changes in the phospholipase activity.

20

Details of functional assays such as those described above are provided in the specific examples, below.

#### Gene Expression

25

30

In another embodiment, test compounds which increase or decrease P2Y<sub>15</sub> GPCR gene expression are identified. A P2Y<sub>15</sub> GPCR polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the P2Y<sub>15</sub> GPCR polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified

as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

5

The level of P2Y15 GPCR mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a P2Y15 GPCR polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labeled amino acids into a P2Y15 GPCR polypeptide.

10

15

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses a P2Y15 GPCR polynucleotide can be used in a cell-based assay system. The P2Y15 GPCR polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

20

#### Pharmaceutical Compositions

The invention also provides pharmaceutical compositions which can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a P2Y15 GPCR polypeptide, P2Y15 GPCR polynucleotide, antibodies which specifically bind to a P2Y15 GPCR polypeptide, or mimetics, agonists, antagonists, or inhibitors of a P2Y15 GPCR polypeptide activity. The compositions can be administered alone or in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, bio-compatible pharmaceutical carrier, including, but not limited to, saline, buffered

25

30

saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

In addition to the active ingredients, these pharmaceutical compositions can contain  
5 suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries  
which facilitate processing of the active compounds into preparations which can be  
used pharmaceutically. Pharmaceutical compositions of the invention can be  
administered by any number of routes including, but not limited to, oral, intravenous,  
intramuscular, intra-arterial, intramedullary, intrapulmonary, intrahepatic, intrathecal,  
10 intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral,  
topical, sublingual, or rectal means. Pharmaceutical compositions for oral  
administration can be formulated using pharmaceutically acceptable carriers well  
known in the art in dosages suitable for oral administration. Such carriers enable the  
pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules,  
15 liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of  
active compounds with solid excipient, optionally grinding a resulting mixture, and  
processing the mixture of granules, after adding suitable auxiliaries, if desired, to  
20 obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers,  
such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn,  
wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxy-  
propylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic  
and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating  
25 or solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone,  
agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores can be used in conjunction with suitable coatings, such as concentrated  
sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone,  
30 carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and  
suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to



the tablets or dragee coatings for product identification or to characterize the quantity of active compound, *i.e.*, dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made  
5 of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid poly-  
10 ethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection  
15 suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes.  
20 Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the  
25 art.

The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or  
30 lyophilizing processes. The pharmaceutical composition can be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric,

sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (Maack Publishing Co., Easton, Pa.). After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

#### Therapeutic indications and methods

This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense nucleic acid molecule, a specific antibody, ribozyme, or a P2Y<sub>15</sub> GPCR polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

A reagent which affects P2Y<sub>15</sub> GPCR activity can be administered to a human cell, either *in vitro* or *in vivo*, to reduce P2Y<sub>15</sub> GPCR activity. The reagent preferably binds to an expression product of a human P2Y<sub>15</sub> GPCR gene. If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells *ex vivo*, an antibody can be added to a preparation of stem cells which have been

- 58 -

removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

5 In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of  
10 targeting to a specific organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its  
15 contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5  $\mu\text{g}$  of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, more preferably about 1.0  $\mu\text{g}$  of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, and even more preferably about 2.0  $\mu\text{g}$  of DNA per 16 nmol of liposome delivered to about  $10^6$  cells. Preferably, a liposome is between about 100 and 500  
20 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

Suitable liposomes for use in the present invention include those liposomes standardly used in, for example, gene delivery methods known to those of skill in the  
25 art. More preferred liposomes include liposomes having a polycationic lipid composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell types, such as a cell-specific ligand exposed on the outer surface of the liposome.

30

Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods which are standard in the art (see, for example, U.S. Patent 5,705,151). Preferably, from about 0.1  $\mu$ g to about 10  $\mu$ g of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5  $\mu$ g to about 5  $\mu$ g of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0  $\mu$ g of polynucleotides is combined with about 8 nmol liposomes.

In another embodiment, antibodies can be delivered to specific tissues *in vivo* using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, Findeis *et al.* *Trends in Biotechnol.* 11, 202-05 (1993); Chiou *et al.*, GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER (J.A. Wolff, ed.) (1994); Wu & Wu, *J. Biol. Chem.* 263, 621-24 (1988); Wu *et al.*, *J. Biol. Chem.* 269, 542-46 (1994); Zenke *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 87, 3655-59 (1990); Wu *et al.*, *J. Biol. Chem.* 266, 338-42 (1991).

#### Determination of a Therapeutically Effective Dose

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases P2Y<sub>15</sub> GPCR activity relative to the P2Y<sub>15</sub> GPCR activity which occurs in the absence of the therapeutically effective dose.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, *e.g.*, ED<sub>50</sub> (the dose therapeutically effective in 50% of the population) and LD<sub>50</sub> (the dose lethal to 50% of the population), can be

- 60 -

determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio,  $LD_{50}/ED_{50}$ .

5      Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage  
10      form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect.  
15      Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on  
20      the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available  
25      to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

30      If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-

established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun," and DEAE- or calcium phosphate-mediated transfection.

Effective *in vivo* dosages of an antibody are in the range of about 5 µg to about 50 µg/kg, about 50 µg to about 5 mg/kg, about 100 µg to about 500 µg/kg of patient body weight, and about 200 to about 250 µg/kg of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective *in vivo* dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1 µg to about 2 mg, about 5 µg to about 500 µg, and about 20 µg to about 100 µg of DNA.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above.

Preferably, a reagent reduces expression of a P2Y<sub>15</sub> GPCR gene or the activity of a P2Y<sub>15</sub> GPCR polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a P2Y<sub>15</sub> GPCR gene or the activity of a P2Y<sub>15</sub> GPCR polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to P2Y<sub>15</sub> GPCR-specific mRNA, quantitative RT-PCR, immunologic detection of a P2Y<sub>15</sub> GPCR polypeptide, or measurement of P2Y<sub>15</sub> GPCR activity.

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can

be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

10

#### Diagnostic Methods

GPCRs also can be used in diagnostic assays for detecting diseases and abnormalities or susceptibility to diseases and abnormalities related to the presence of mutations in the nucleic acid sequences which encode a GPCR. Such diseases, by way of example, are related to cell transformation, such as tumors and cancers, and various cardiovascular disorders, including hypertension and hypotension, as well as diseases arising from abnormal blood flow, abnormal angiotensin-induced aldosterone secretion, and other abnormal control of fluid and electrolyte homeostasis.

15

According to the present invention, differences can be determined between the cDNA or genomic sequence encoding a P2Y<sub>15</sub> GPCR in individuals afflicted with a disease and in normal individuals. If a mutation is observed in some or all of the afflicted individuals but not in normal individuals, then the mutation is likely to be the causative agent of the disease.

20

Sequence differences between a reference gene and a gene having mutations can be revealed by the direct DNA sequencing method. In addition, cloned DNA segments can be employed as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequencing primer can be used with a double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by

25

30

conventional procedures using radiolabeled nucleotides or by automatic sequencing procedures using fluorescent tags.

Genetic testing based on DNA sequence differences can be carried out by detection  
5 of alteration in electrophoretic mobility of DNA fragments in gels with or without  
denaturing agents. Small sequence deletions and insertions can be visualized, for  
example, by high resolution gel electrophoresis. DNA fragments of different  
sequences can be distinguished on denaturing formamide gradient gels in which the  
mobilities of different DNA fragments are retarded in the gel at different positions  
10 according to their specific melting or partial melting temperatures (*see, e.g., Myers et al., Science 230, 1242, 1985*). Sequence changes at specific locations can also be  
revealed by nuclease protection assays, such as RNase and S 1 protection or the  
chemical cleavage method (*e.g., Cotton et al., Proc. Natl. Acad. Sci. USA 85, 4397-4401, 1985*). Thus, the detection of a specific DNA sequence can be performed  
15 by methods such as hybridization, RNase protection, chemical cleavage, direct DNA  
sequencing or the use of restriction enzymes and Southern blotting of genomic DNA.  
In addition to direct methods such as gel-electrophoresis and DNA sequencing,  
mutations can also be detected by *in situ* analysis.

20 Altered levels of a P2Y<sub>15</sub> GPCR also can be detected in various tissues. Assays  
used to detect levels of the receptor polypeptides in a body sample, such as blood or a  
tissue biopsy, derived from a host are well known to those of skill in the art and  
include radioimmunoassays, competitive binding assays, Western blot analysis, and  
ELISA assays.

25 All patents and patent applications cited in this disclosure are expressly incorporated  
herein by reference. The above disclosure generally describes the present invention.  
A more complete understanding can be obtained by reference to the following  
specific examples which are provided for purposes of illustration only and are not  
30 intended to limit the scope of the invention.



**EXAMPLE 1***Detection of P2Y15 GPCR activity*

The polynucleotide of SEQ ID NO: 1, 3, or 5 is inserted into the expression vector pCEV4 and the expression vector pCEV4-P2Y15 GPCR polypeptide obtained is transfected into human embryonic kidney 293 cells. These cells are scraped from a culture flask into 5 ml of Tris HCl, 5 mM EDTA, pH 7.5, and lysed by sonication. Cell lysates are centrifuged at 1000 rpm for 5 minutes at 4°C. The supernatant is centrifuged at 30,000 x g for 20 minutes at 4°C. The pellet is suspended in binding buffer containing 50 mM Tris HCl, 5 mM MgSO<sub>4</sub>, 1 mM EDTA, 100 mM NaCl, pH 7.5, supplemented with 0.1 % BSA, 2 µg/ml aprotinin, 0.5 mg/ml leupeptin, and 10 µg/ml phosphoramidon. Optimal membrane suspension dilutions, defined as the protein concentration required to bind less than 10 % of the added radioligand, are added to 96-well polypropylene microtiter plates containing <sup>125</sup>I-labeled ligand or test compound, non-labeled peptides, and binding buffer to a final volume of 250 µl.

In equilibrium saturation binding assays, membrane preparations are incubated in the presence of increasing concentrations (0.1 nM to 4 nM) of <sup>125</sup>I-labeled ligand or test compound (specific activity 2200 Ci/mmol). The binding affinities of different test compounds are determined in equilibrium competition binding assays, using 0.1 nM <sup>125</sup>I-peptide in the presence of twelve different concentrations of each test compound.

Binding reaction mixtures are incubated for one hour at 30°C. The reaction is stopped by filtration through GF/B filters treated with 0.5% polyethyleneimine, using a cell harvester. Radioactivity is measured by scintillation counting, and data are analyzed by a computerized non-linear regression program.

Non-specific binding is defined as the amount of radioactivity remaining after incubation of membrane protein in the presence of 100 nM of unlabeled peptide. Protein concentration is measured by the Bradford method using Bio-Rad Reagent,

- 65 -

with bovine serum albumin as a standard. It is shown that the polypeptide of SEQ ID NO: 2 has a P2Y<sub>15</sub> GPCR activity.

## **EXAMPLE 2**

### 5     *Expression of recombinant human P2Y<sub>15</sub> GPCR*

The *Pichia pastoris* expression vector pPICZB (Invitrogen, San Diego, CA) is used to produce large quantities of a human P2Y<sub>15</sub> GPCR polypeptides in yeast. The human P2Y<sub>15</sub> GPCR polypeptide-encoding DNA sequence is derived from the  
10     nucleotide sequence shown in SEQ ID NO:1. Before insertion into vector pPICZB the DNA sequence is modified by well known methods in such a way that it contains at its 5'-end an initiation codon and at its 3'-end an enterokinase cleavage site, a His6 reporter tag and a termination codon. Moreover, at both termini recognition  
15     sequences for restriction endonucleases are added and after digestion of the multiple cloning site of pPICZ B with the corresponding restriction enzymes the modified polypeptide encoding DNA sequence is ligated into pPICZB. This expression vector is designed for inducible expression in *Pichia pastoris*, expression is driven by a yeast promoter. The resulting pPICZ/md-His6 vector is used to transform the yeast.

20     The yeast are cultivated under usual conditions in 5 liter shake flasks and the recombinantly produced protein isolated from the culture by affinity chromatography (Ni-NTA-Resin) in the presence of 8 M urea. The bound polypeptide is eluted with buffer, pH 3.5, and neutralized. Separation of the P2Y<sub>15</sub> GPCR polypeptide from the His6 reporter tag is accomplished by site-specific proteolysis using enterokinase  
25     (Invitrogen, San Diego, CA) according to manufacturer's instructions. Purified human P2Y<sub>15</sub> GPCR polypeptide is obtained.

**EXAMPLE 3***Radioligand binding assays*

Human embryonic kidney 293 cells transfected with a polynucleotide which expresses human P2Y<sub>15</sub> GPCR are scraped from a culture flask into 5 ml of Tris HCl, 5 mM EDTA, pH 7.5, and lysed by sonication. Cell lysates are centrifuged at 1000 rpm for 5 minutes at 4°C. The supernatant is centrifuged at 30,000 x g for 20 minutes at 4°C. The pellet is suspended in binding buffer containing 50 mM Tris HCl, 5 mM MgSO<sub>4</sub>, 1 mM EDTA, 100 mM NaCl, pH 7.5, supplemented with 0.1 % BSA, 2 µg/ml aprotinin, 0.5 mg/ml leupeptin, and 10 µg/ml phosphoramidon. Optimal membrane suspension dilutions, defined as the protein concentration required to bind less than 10 % of the added radioligand, are added to 96-well polypropylene microtiter plates containing <sup>125</sup>I-labeled ligand or test compound, non-labeled peptides, and binding buffer to a final volume of 250 µl.

In equilibrium saturation binding assays, membrane preparations are incubated in the presence of increasing concentrations (0.1 nM to 4 nM) of <sup>125</sup>I-labeled ligand or test compound (specific activity 2200 Ci/mmol). The binding affinities of different test compounds are determined in equilibrium competition binding assays, using 0.1 nM <sup>125</sup>I-peptide in the presence of twelve different concentrations of each test compound.

Binding reaction mixtures are incubated for one hour at 30 °C. The reaction is stopped by filtration through GF/B filters treated with 0.5% polyethyleneimine, using a cell harvester. Radioactivity is measured by scintillation counting, and data are analyzed by a computerized non-linear regression program.

Non-specific binding is defined as the amount of radioactivity remaining after incubation of membrane protein in the presence of 100 nM of unlabeled peptide. Protein concentration is measured by the Bradford method using Bio-Rad Reagent, with bovine serum albumin as a standard. A test compound which increases the radioactivity of membrane protein by at least 15% relative to radioactivity of

membrane protein which was not incubated with a test compound is identified as a compound which binds to a human P2Y<sub>15</sub> GPCR polypeptide.

#### **EXAMPLE 4**

##### 5 *Effect of a test compound on human P2Y<sub>15</sub> GPCR-mediated cyclic AMP formation*

Receptor-mediated inhibition of cAMP formation can be assayed in host cells which express human P2Y<sub>15</sub> GPCR. Cells are plated in 96-well plates and incubated in Dulbecco's phosphate buffered saline (PBS) supplemented with 10 mM HEPES,  
10 5 mM theophylline, 2 µg/ml aprotinin, 0.5 mg/ml leupeptin, and 10 µg/ml phosphoramidon for 20 minutes at 37°C in 5% CO<sub>2</sub>. A test compound is added and incubated for an additional 10 minutes at 37°C. The medium is aspirated, and the reaction is stopped by the addition of 100 mM HCl. The plates are stored at 4°C for 15 minutes. cAMP content in the stopping solution is measured by radioimmunoassay.

15

Radioactivity is quantified using a gamma counter equipped with data reduction software. A test compound which decreases radioactivity of the contents of a well relative to radioactivity of the contents of a well in the absence of the test compound is identified as a potential inhibitor of cAMP formation. A test compound which  
20 increases radioactivity of the contents of a well relative to radioactivity of the contents of a well in the absence of the test compound is identified as a potential enhancer of cAMP formation.

#### **EXAMPLE 5**

##### 25 *Effect of a test compound on the mobilization of intracellular calcium*

Intracellular free calcium concentration can be measured by microspectrofluorometry using the fluorescent indicator dye Fura-2/AM (Bush *et al.*, *J. Neurochem.* 57, 562-74, 1991). Stably transfected cells are seeded onto a 35 mm culture dish  
30 containing a glass coverslip insert. Cells are washed with HBS, incubated with a test compound, and loaded with 100 µl of Fura-2/AM (10 µM) for 20-40 minutes. After

- 68 -

washing with HBS to remove the Fura-2/AM solution, cells are equilibrated in HBS for 10-20 minutes. Cells are then visualized under the 40X objective of a Leitz Fluovert FS microscope.

5 Fluorescence emission is determined at 510 nm, with excitation wavelengths alternating between 340 nm and 380 nm. Raw fluorescence data are converted to calcium concentrations using standard calcium concentration curves and software analysis techniques. A test compound which increases the fluorescence by at least 15% relative to fluorescence in the absence of a test compound is identified as a  
10 compound which mobilizes intracellular calcium.

#### **EXAMPLE 6**

##### *Effect of a test compound on phosphoinositide metabolism*

15 Cells which stably express human P2Y<sub>15</sub> GPCR cDNA are plated in 96-well plates and grown to confluence. The day before the assay, the growth medium is changed to 100 µl of medium containing 1% serum and 0.5 µCi <sup>3</sup>H-myo-inositol. The plates are incubated overnight in a CO<sub>2</sub> incubator (5% CO<sub>2</sub> at 37°C). Immediately before the assay, the medium is removed and replaced by 200 µl of PBS containing 10 mM  
20 LiCl, and the cells are equilibrated with the new medium for 20 minutes. During this interval, cells also are equilibrated with antagonist, added as a 10 µl aliquot of a 20-fold concentrated solution in PBS.

The <sup>3</sup>H-inositol phosphate accumulation from inositol phospholipid metabolism is  
25 started by adding 10 µl of a solution containing a test compound. To the first well 10 µl are added to measure basal accumulation. Eleven different concentrations of test compound are assayed in the following 11 wells of each plate row. All assays are performed in duplicate by repeating the same additions in two consecutive plate rows.

30

The plates are incubated in a CO<sub>2</sub> incubator for one hour. The reaction is terminated by adding 15 µl of 50% v/v trichloroacetic acid (TCA), followed by a 40 minute incubation at 4 °C. After neutralizing TCA with 40 µl of 1 M Tris, the content of the wells is transferred to a Multiscreen HV filter plate (Millipore) containing Dowex AG1-X8 (200-400 mesh, formate form). The filter plates are prepared by adding 200 µl of Dowex AG1-X8 suspension (50% v/v, water:resin) to each well. The filter plates are placed on a vacuum manifold to wash or elute the resin bed. Each well is washed 2 times with 200 µl of water, followed by 2 x 200 µl of 5 mM sodium tetraborate/60 mM ammonium formate.

10.

The <sup>3</sup>H-IPs are eluted into empty 96-well plates with 200 µl of 1.2 M ammonium formate/0.1 formic acid. The content of the wells is added to 3 ml of scintillation cocktail, and radioactivity is determined by liquid scintillation counting.

15

### EXAMPLE 7

#### *Receptor Binding Methods*

20

Standard Binding Assays. Binding assays are carried out in a binding buffer containing 50 mM HEPES, pH 7.4, 0.5% BSA, and 5 mM MgCl<sub>2</sub>. The standard assay for radioligand (e.g., <sup>125</sup>I- test compound) binding to membrane fragments comprising P2Y<sub>15</sub> GPCR polypeptides is carried out as follows in 96 well microtiter plates (e.g., Dynatech Immulon II Removawell plates). Radioligand is diluted in binding buffer+ PMSF/Baci to the desired cpm per 50 µl, then 50 µl aliquots are added to the wells. For non-specific binding samples, 5 µl of 40 µM cold ligand also is added per well. Binding is initiated by adding 150 µl per well of membrane diluted to the desired concentration (10-30 µg membrane protein/well) in binding buffer+ PMSF/Baci. Plates are then covered with Linbro mylar plate sealers (Flow Labs) and placed on a Dynatech Microshaker II. Binding is allowed to proceed at room temperature for 1-2 hours and is stopped by centrifuging the plate for 15 minutes at 2,000 x g. The supernatants are decanted, and the membrane pellets

30

- 70 -

are washed once by addition of 200  $\mu$ l of ice cold binding buffer, brief shaking, and recentrifugation. The individual wells are placed in 12 x 75 mm tubes and counted in an LKB Gammamaster counter (78% efficiency). Specific binding by this method is identical to that measured when free ligand is removed by rapid (3-5 seconds) filtration and washing on polyethyleneimine-coated glass fiber filters.

Three variations of the standard binding assay are also used.

1. Competitive radioligand binding assays with a concentration range of cold ligand vs.  $^{125}$ I-labeled ligand are carried out as described above with one modification. All dilutions of ligands being assayed are made in 40X PMSF/Baci to a concentration 40X the final concentration in the assay. Samples of peptide (5  $\mu$ l each) are then added per microtiter well. Membranes and radioligand are diluted in binding buffer without protease inhibitors. Radioligand is added and mixed with cold ligand, and then binding is initiated by addition of membranes.
2. Chemical cross-linking of radioligand with receptor is done after a binding step identical to the standard assay. However, the wash step is done with binding buffer minus BSA to reduce the possibility of non-specific cross-linking of radioligand with BSA. The cross-linking step is carried out as described below.
3. Larger scale binding assays to obtain membrane pellets for studies on solubilization of receptor:ligand complex and for receptor purification are also carried out. These are identical to the standard assays except that (a) binding is carried out in polypropylene tubes in volumes from 1-250 ml, (b) concentration of membrane protein is always 0.5 mg/ml, and (c) for receptor purification, BSA concentration in the binding buffer is reduced to 0.25%, and the wash step is done with binding buffer without BSA, which reduces BSA contamination of the purified receptor.

**EXAMPLE 8***Chemical Cross-Linking of Radioligand to Receptor*

5 After a radioligand binding step as described above, membrane pellets are resuspended in 200  $\mu$ l per microtiter plate well of ice-cold binding buffer without BSA. Then 5  $\mu$ l per well of 4 mM N-5-azido-2-nitrobenzoyloxysuccinimide (ANB-NOS, Pierce) in DMSO is added and mixed. The samples are held on ice and UV-irradiated for 10 minutes with a Mineralight R-52G lamp (UVP Inc., San  
10 Gabriel, Calif.) at a distance of 5-10 cm. Then the samples are transferred to Eppendorf microfuge tubes, the membranes pelleted by centrifugation, supernatants removed, and membranes solubilized in Laemmli SDS sample buffer for polyacrylamide gel electrophoresis (PAGE). PAGE is carried out as described below. Radiolabeled proteins are visualized by autoradiography of the dried gels with Kodak  
15 XAR film and DuPont image intensifier screens.

**EXAMPLE 9***Membrane Solubilization*

20 Membrane solubilization is carried out in buffer containing 25 mM Tris, pH 8, 10% glycerol (w/v) and 0.2 mM  $\text{CaCl}_2$  (solubilization buffer). The highly soluble detergents including Triton X-100, deoxycholate, deoxycholate:lysolecithin, CHAPS, and zwittergent are made up in solubilization buffer at 10% concentrations and stored as frozen aliquots. Lysolecithin is made up fresh because of insolubility upon  
25 freeze-thawing and digitonin is made fresh at lower concentrations due to its more limited solubility.

To solubilize membranes, washed pellets after the binding step are resuspended free of visible particles by pipetting and vortexing in solubilization buffer at 100,000 x g  
30 for 30 minutes. The supernatants are removed and held on ice and the pellets are discarded.



**EXAMPLE 10***Assay of Solubilized Receptors*

5 After binding of  $^{125}\text{I}$  ligands and solubilization of the membranes with detergent, the intact R:L complex can be assayed by four different methods. All are carried out on ice or in a cold room at 4-10°C).

10 Column chromatography (Knuhtsen *et al.*, *Biochem. J.* 254, 641-647, 1988).  
Sephadex G-50 columns (8 x 250 mm) are equilibrated with solubilization buffer containing detergent at the concentration used to solubilize membranes and 1 mg/ml bovine serum albumin. Samples of solubilized membranes (0.2-0.5 ml) are applied to the columns and eluted at a flow rate of about 0.7 ml/minute. Samples (0.18 ml) are collected. Radioactivity is determined in a gamma counter. Void volumes of the  
15 columns are determined by the elution volume of blue dextran. Radioactivity eluting in the void volume is considered bound to protein. Radioactivity eluting later, at the same volume as free  $^{125}\text{I}$  ligands, is considered non-bound.

20 Polyethyleneglycol precipitation (Cuatrecasas, *Proc. Natl. Acad. Sci. USA* 69, 318-322, 1972). For a 100  $\mu\text{l}$  sample of solubilized membranes in a 12 x 75 mm polypropylene tube, 0.5 ml of 1% (w/v) bovine gamma globulin (Sigma) in 0.1 M sodium phosphate buffer is added, followed by 0.5 ml of 25% (w/v) polyethyleneglycol (Sigma) and mixing. The mixture is held on ice for 15 minutes. Then 3 ml of 0.1 M sodium phosphate, pH 7.4, is added per sample. The samples are  
25 rapidly (1-3 seconds) filtered over Whatman GF/B glass fiber filters and washed with 4 ml of the phosphate buffer. PEG-precipitated receptor :  $^{125}\text{I}$ -ligand complex is determined by gamma counting of the filters.

30 GFB/PEI filter binding (Bruns *et al.*, *Analytical Biochem.* 132, 74-81, 1983).  
Whatman GF/B glass fiber filters are soaked in 0.3% polyethyleneimine (PEI, Sigma) for 3 hours. Samples of solubilized membranes (25-100  $\mu\text{l}$ ) are replaced in 12 x

- 73 -

75 mm polypropylene tubes. Then 4 ml of solubilization buffer without detergent is added per sample and the samples are immediately filtered through the GFB/PEI filters (1-3 seconds) and washed with 4 ml of solubilization buffer. CPM of receptor :  $^{125}$ I-ligand complex adsorbed to filters are determined by gamma counting.

5

Charcoal/Dextran (Paul and Said, *Peptides* 7[Suppl. 1],147-149, 1986). Dextran T70 (0.5 g, Pharmacia) is dissolved in 1 liter of water, then 5 g of activated charcoal (Norit A, alkaline; Fisher Scientific) is added. The suspension is stirred for 10 minutes at room temperature and then stored at 4 °C. until use. To measure R:L complex, 4 parts by volume of charcoal/dextran suspension are added to 1 part by volume of solubilized membrane. The samples are mixed and held on ice for 2 minutes and then centrifuged for 2 minutes at 11,000 x g in a Beckman microfuge. Free radioligand is adsorbed charcoal/dextran and is discarded with the pellet. Receptor :  $^{125}$ I-ligand complexes remain in the supernatant and are determined by gamma counting.

15

### EXAMPLE 11

#### *Receptor Purification*

Binding of biotinyl-receptor to GH<sub>4</sub> C1 membranes is carried out as described above. Incubations are for 1 hour at room temperature. In the standard purification protocol, the binding incubations contain 10 nM Bio-S29.  $^{125}$ I ligand is added as a tracer at levels of 5,000-100,000 cpm per mg of membrane protein. Control incubations contain 10  $\mu$ M cold ligand to saturate the receptor with non-biotinylated ligand.

25

Solubilization of receptor:ligand complex also is carried out as described above, with 0.15% deoxycholate:lysolecithin in solubilization buffer containing 0.2 mM MgCl<sub>2</sub>, to obtain 100,000 x g supernatants containing solubilized R:L complex.

30

Immobilized streptavidin (streptavidin cross-linked to 6% beaded agarose, Pierce Chemical Co.; "SA-agarose") is washed in solubilization buffer and added to the

solubilized membranes as 1/30 of the final volume. This mixture is incubated with constant stirring by end-over-end rotation for 4-5 hours at 4-10 °C. Then the mixture is applied to a column and the non-bound material is washed through. Binding of radioligand to SA-agarose is determined by comparing cpm in the 100,000 x g  
5 supernatant with that in the column effluent after adsorption to SA-agarose. Finally, the column is washed with 12-15 column volumes of solubilization buffer+0.15% deoxycholate:lysolecithin +1/500 (vol/vol) 100 x 4pase.

The streptavidin column is eluted with solubilization buffer+0.1 mM EDTA+0.1 mM  
10 EGTA+0.1 mM GTP-gamma-S (Sigma)+0.15% (wt/vol) deoxycholate:lysolecithin +1/1000 (vol/vol) 100.times.4pase. First, one column volume of elution buffer is passed through the column and flow is stopped for 20-30 minutes. Then 3-4 more column volumes of elution buffer are passed through. All the eluates are pooled.

15 Eluates from the streptavidin column are incubated overnight (12-15 hours) with immobilized wheat germ agglutinin (WGA agarose, Vector Labs) to adsorb the receptor via interaction of covalently bound carbohydrate with the WGA lectin. The ratio (vol/vol) of WGA-agarose to streptavidin column eluate is generally 1:400. A range from 1:1000 to 1:200 also can be used. After the binding step, the resin is  
20 pelleted by centrifugation, the supernatant is removed and saved, and the resin is washed 3 times (about 2 minutes each) in buffer containing 50 mM HEPES, pH 8, 5 mM MgCl<sub>2</sub>, and 0.15% deoxycholate:lysolecithin. To elute the WGA-bound receptor, the resin is extracted three times by repeated mixing (vortex mixer on low speed) over a 15-30 minute period on ice, with 3 resin columns each time, of 10 mM  
25 N-N'-N''-triacetylchitotriose in the same HEPES buffer used to wash the resin. After each elution step, the resin is centrifuged down and the supernatant is carefully removed, free of WGA-agarose pellets. The three, pooled eluates contain the final, purified receptor. The material non-bound to WGA contain G protein subunits specifically eluted from the streptavidin column, as well as non-specific con-  
30 taminants. All these fractions are stored frozen at -90°C.

**EXAMPLE 12***Identification of test compounds that bind to P2Y15 GPCR polypeptides*

Purified P2Y15 GPCR polypeptides comprising a glutathione-S-transferase protein  
5 and absorbed onto glutathione-derivatized wells of 96-well microtiter plates are  
contacted with test compounds from a small molecule library at pH 7.0 in a physio-  
logical buffer solution. P2Y15 GPCR polypeptides comprise an amino acid  
sequence shown in SEQ ID NO:2. The test compounds comprise a fluorescent tag.  
The samples are incubated for 5 minutes to one hour. Control samples are incubated  
10 in the absence of a test compound.

The buffer solution containing the test compounds is washed from the wells.  
Binding of a test compound to a P2Y15 GPCR polypeptide is detected by  
fluorescence measurements of the contents of the wells. A test compound which  
15 increases the fluorescence in a well by at least 15% relative to fluorescence of a well  
in which a test compound was not incubated is identified as a compound which binds  
to a P2Y15 GPCR polypeptide.

**EXAMPLE 13***Identification of a test compound which decreases P2Y15 GPCR gene expression*

A test compound is administered to a culture of human gastric cells and incubated at  
37°C for 10 to 45 minutes. A culture of the same type of cells incubated for the same  
time without the test compound provides a negative control.

25 RNA is isolated from the two cultures as described in Chirgwin *et al.*, *Biochem. 18*,  
5294-99, 1979). Northern blots are prepared using 20 to 30 µg total RNA and  
hybridized with a <sup>32</sup>P-labeled P2Y15 GPCR-specific probe at 65°C in Express-hyb  
(CLONTECH). The probe comprises at least 11 contiguous nucleotides selected  
30 from the complement of SEQ ID NO:1. A test compound which decreases the

- 76 -

P2Y15 GPCR-specific signal relative to the signal obtained in the absence of the test compound is identified as an inhibitor of P2Y15 GPCR gene expression.

#### **EXAMPLE 14**

5     *Treatment of a disease in which human P2Y15 GPCR is overexpressed with a reagent which specifically binds to a P2Y15 GPCR gene product*

Synthesis of antisense P2Y15 GPCR oligonucleotides comprising at least 11 contiguous nucleotides selected from the complement of SEQ ID NO:1 is performed  
10     on a Pharmacia Gene Assembler series synthesizer using the phosphoramidite procedure (Uhlmann *et al.*, *Chem. Rev.* 90, 534-83, 1990). Following assembly and deprotection, oligonucleotides are ethanol-precipitated twice, dried, and suspended in phosphate-buffered saline (PBS) at the desired concentration. Purity of these oligonucleotides is tested by capillary gel electrophoreses and ion exchange HPLC.  
15     Endotoxin levels in the oligonucleotide preparation are determined using the Luminous Amebocyte Assay (Bang, *Biol. Bull. (Woods Hole, Mass.)* 105, 361-362, 1953).

The antisense oligonucleotides are administered to a patient. The severity of the  
20     patient's disease is decreased.

#### **EXAMPLE 15**

*Tissue-specific expression of P2Y15 GPCR*

25     In an analysis of P2Y15 GPCR gene expression, 25 µg of total RNA from the following sources were used as template in reactions to synthesize first-strand cDNA for expression profiling : Human Total RNA Panel I-V (Clontech Laboratories, Palo Alto, CA, USA), normal human lung primary cell lines (BioWhittaker Clonetics, Walkersville, MD, USA), human umbilical vein endothelial cells (HUVECs)  
30     (Kurabo, Osaka, Japan), several common cell lines (ATCC, Washington, DC), and various cells purified from peripheral blood. First-strand cDNA was synthesized

- 77 -

using oligo (dT) (Nippon Gene Research Laboratories, Sendai, Japan) and the SUPERScript™ First-Strand Synthesis System for RT-PCR (Life Technologies, Rockville, MD) according to the manufacturer's protocol. For these samples, 1/1250<sup>th</sup> of the synthesized first-strand cDNA was subsequently used as template for  
5 quantitative PCR. Additional samples were purchased as presynthesized cDNAs (Human Immune System MTC Panel and Human Blood Fractions MTC Panel, Clontech Laboratories), and for these, 10 ng of cDNA was used as template for quantitative PCR.

10 Quantitative PCR was performed in a LightCycler (Roche Molecular Biochemicals, Indianapolis, IN) with oligonucleotide primers 5'-TTCGGATCGAATCTCGCCTGCT-3' (SEQ ID NO:7) and 5'-TGCTTGCTCAAGGTTCCCGCTTA-3' (SEQ ID NO:8) in the presence of the DNA-binding fluorescent dye SYBR Green I. Results were then converted into copy numbers of the gene transcript per ng of template cDNA by fitting to  
15 a standard curve. The standard curve was derived by simultaneously performing the quantitative PCR reaction on PCR products of known concentrations amplified beforehand from the target gene.

To correct for differences in mRNA transcription levels per cell in the various tissue  
20 types, a normalization procedure was performed using similarly calculated expression levels of five different housekeeping genes: glyceraldehyde-3-phosphatase (G3PDH), hypoxanthine guanine phosphoribosyl transferase (HPRT), beta-actin, porphobilinogen deaminase (PBGD), and beta-2-microglobulin. The level of house-keeping gene expression is considered to be relatively constant for all tissues (Adams  
25 et al., 1993, Adams et al., 1995, Liew et al., 1994) and therefore can be used as a gauge to approximate relative numbers of cells per ng of cDNA template. Expression levels of the five housekeeping genes in all tissue samples were measured in three independent reactions per gene using the LightCycler and a constant amount (25 µg) of starting RNA. The calculated copy numbers for each gene, derived from com-  
30 parison with simultaneously reacted standards of known concentrations, were recorded and converted into a percentage of the sum of the copy numbers of the gene

in all tissue samples. For each tissue sample, the sum of the percentage values for each gene was calculated, and a normalization factor was calculated by dividing the sum percentage value for each tissue by the sum percentage value of one of the tissues arbitrarily selected as a standard. To normalize an experimentally obtained value for the expression of a particular gene in a tissue sample, the obtained value was multiplied by the normalization factor for the tissue tested. This normalization method was used for all tissues except those derived from the Human Blood Fractions MTC Panel, which were normalized against the single housekeeping gene, beta-2-microglobulin, due to wide variation in other housekeeping gene expression in these tissues depending on activation status. The results of this expression profiling are given in Fig. 1, showing the normalized values for the copy numbers of mRNA per 10 ng of first-strand cDNA in each sample tested.

#### **EXAMPLE 16**

##### ***Determination of the P2Y15 GPCR ligand***

The polynucleotide of SEQ ID NO: 1 was cloned using human genomic DNA as a template and performing PCR with primers 5'-GCCAAACTGAACTCTCTTGTTCCTTGC-3' (SEQ ID NO:9) and 5'-GCCCTGGCTTTGGCACATGATTAC-3' (SEQ ID NO:10) and a blend of HotStarTaq (Qiagen, Hilden, Germany) and Pfu Turbo (Stratagene, La Jolla, CA) polymerases. The PCR products were cloned into pCRII-TOPO (Invitrogen, Carlsbad, CA), cycle-sequenced with an ABI Prism Dye Terminator Cycle Sequencing Reaction Kit (Applied Biosystems, Foster City, CA), and analyzed on an ABI Prism 377 sequencing system (Applied Biosystems). The cDNA was then subcloned into a modified pDisplay vector (Invitrogen) to append an N-terminal HA epitope and Ig signal sequence. The expression vector pDisplay-P2Y15 GPCR construct obtained was transfected into human embryonic kidney 293 cells using Lipofectamine (Invitrogen). Expression on the cell surface was verified by staining the cells with phycoerythrin-labeled anti-HA antibody (Santa Cruz Biotechnology, Santa Cruz, CA) and measuring cell-bound fluorescence on a FACSsort (Becton-Dickinson, Franklin Lakes, NJ). Stably transfected clones were

- 79 -

then generated by limiting dilution followed by selection in G418. Finally, cell-surface expression of the P2Y15 GPCR polypeptide in the clones was reconfirmed by FACS.

Ligand screening was performed in a  $\text{Ca}^{2+}$  flux assay as follows. Stably transfected  
5 P2Y15 GPCR-expressing cells were seeded into 96-well plates and incubated  
overnight at 37°C. in a tissue culture incubator. The culture medium was aspirated  
and replaced with 100  $\mu$ l of loading buffer containing 0.1% BSA, 20 mM HEPES,  
1 mM probenecid, 0.01% pluronic F127, and 1  $\mu$ M Fluo-3-AM (Molecular Probes,  
Eugene, OR) in HBSS, and incubated for 1 hour at room temperature. The cells were  
10 then washed gently 3 times with wash buffer containing 0.1% BSA, 20 mM HEPES,  
1 mM probenecid in HBSS. The washed cells were placed in an FDSS6000  
functional drug screening system (Hamamatsu Photonics, Hamamatsu, Japan) and  
changes in cellular fluorescence were measured after adding serial dilutions of  
potential ligands. A panel of about 130 potential ligands was assembled by selecting  
15 known ligands of the GPCRs most closely related to P2Y15 GPCR as well as several  
naturally occurring chemical relatives of the ligands. The library included various  
bioactive lipids, eicosanoids, peptides, cannabinoids, chemokines, nucleosides,  
nucleotides and chemically related substances. The ligands were generally  
purchased from either Sigma or R&D Systems.

20 Among the potential ligands tested, only AMP and adenosine were able to induce a  
response in the transfectants while not inducing a similar response in either  
nontransfected HEK293 cells or HEK293 cells stably transfected with the control  
orphan GPCR P2Y8 in an identical vector construct. We detected a calcium response  
25 with an EC<sub>50</sub> of 920 nM for AMP and 670 nM for adenosine (Fig. 2A). Both stable  
transfectants and nontransfected cells mobilized calcium in response to ATP, AMP  
OR ADENOSINE RECEPTOR LIGAND, and UTP, consistent with previous reports  
of HEK293 endogenously expressing P2Y1 and P2Y2 receptors (Schachter, J. B.,  
Sromek, S. M., Nicholas, R. A., and Harden, T. K. (1997) *Neuropharmacology* 36,  
30 1181-1187). Further analysis by RT-PCR showed that the nucleotide receptors P2Y4,  
P2Y12, and P2Y13 and the adenosine receptors A2A and A2B (previously reported



in (Sunahara, R. K., Dessauer, C. W., and Gilman, A. G. (1996) *Annu Rev Pharmacol Toxicol* 36, 461-480)) are also expressed in HEK293 cells. Despite the endogenous expression of the adenosine receptors, calcium mobilization responses to adenosine in nontransfected cells could only be detected at very high adenosine concentrations, and showed only a very weak response.

### EXAMPLE 17

#### *Cyclic AMP production assay*

To determine the effect of P2Y<sub>15</sub> stimulation on adenylyl cyclase activity, cyclic AMP accumulation in P2Y<sub>15</sub>-HEK293 stable transfectants in response to AMP and adenosine was measured with the Tropix cAMP screen (Applied Biosystems) according to the manufacturer's protocol. Briefly, stable transfectants and control cells ( $1 \times 10^5$  cells/well) were cultured for two hours with or without 1  $\mu$ M pertussis toxin, then treated for 30 min with 10  $\mu$ M forskolin and serial dilutions of AMP or adenosine. The cells were then lysed and the cAMP produced was measured by a cAMP-specific ELISA. Concentrations of cAMP produced were calculated by comparing against cAMP standards measured simultaneously. Stimulation with either ligand alone gave only minor responses barely above the detection limit. In the presence of 10  $\mu$ M forskolin, however, both AMP and adenosine induced the generation of cyclic AMP in a dose dependent manner, with an EC<sub>50</sub> of 214 nM for AMP and 327 nM for adenosine (Fig. 2B). Nontransfected HEK293 cells similarly generated cyclic AMP in response to adenosine, likely due to the stimulation of endogenously expressed adenosine receptors, but did not respond strongly to AMP. The production of cyclic AMP in response to either ligand was not affected by pretreatment of the cells for two hours with 1  $\mu$ M pertussis toxin, indicating that P2Y<sub>15</sub> is coupled to an adenylate cyclase-stimulating G protein.

**EXAMPLE 18***Receptor binding assay*

10<sup>5</sup> cells per well in 96-well plates were washed twice for 1 h with DMEM medium. Then wheatgerm agglutinin SPA beads (Amersham) at 1 mg/well were added, followed 1 h later by increasing concentrations of [<sup>3</sup>H]-Adenosine or [<sup>3</sup>H]-AMP (Amersham), in a constant volume of HBS: 10 mM Hepes, 130 mM NaCl, 5 mM KCl, 1 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub> and 1 g/l glucose (6 µl/well). After incubating at 4°C for 16 h, the plates were centrifuged for 10 min at 1500 rpm and then scintillation measured on a TopCount automated scintillation counter. Binding of [<sup>32</sup>P]-AMP was carried out under the same conditions except that instead of using SPA beads, at the end of the incubation, cells were washed three times by vacuum filtration and 100 µl scintillation fluid was added to the wells. Non-specific binding measurements were carried out under the same conditions but with either an excess of cold ligand (2.5 mM). K<sub>d</sub> values were determined by non-linear regression using the program Prism (Graph Pad).

Saturation binding analysis of the ligands to the P2Y<sub>15</sub> receptor in stable transfectants gave K<sub>d</sub> values of 12.0 µM for <sup>3</sup>H-adenosine (Fig. 3A) and 18.6 µM for <sup>3</sup>H-AMP (data not shown). Since AMP can be dephosphorylated to adenosine by ectonucleotidases, we repeated the binding analysis with adenosine 5'-[<sup>32</sup>P] monophosphate to confirm that the binding being measured was AMP and not adenosine. This resulted in a similar binding curve with a K<sub>d</sub> of 18.8 µM (Fig. 3B), indicating that AMP itself, and not a breakdown product, was binding to the receptor.

**EXAMPLE 19***Competitive binding assay*

Competitive binding experiments were carried out as follows: 10<sup>5</sup> cells per well in 96-well plates were washed twice for 1 h with DMEM medium. Then wheatgerm agglutinin SPA beads (Amersham) at 1 mg/well were added, followed 1 h later by

10  $\mu\text{M}$  [ $^3\text{H}$ ]-Adenosine or 10  $\mu\text{M}$  [ $^3\text{H}$ ]-AMP (Amersham), and increasing concentrations of unlabeled ligand in a constant volume of HBS: 10 mM Hepes, 130 mM NaCl, 5 mM KCl, 1 mM  $\text{MgCl}_2$ , 1 mM  $\text{CaCl}_2$  and 1 g/l glucose (6  $\mu\text{l}$ /well). After incubating at 4 °C for 16 h, the plates were centrifuged for 10 min at 1500 rpm and then scintillation measured on a TopCount automated scintillation counter.  $K_i$  values were determined by non-linear regression using the program Prism (Graph Pad).

Unlabeled adenosine was able to block the binding of 10  $\mu\text{M}$   $^3\text{H}$ -AMP to transfectants ( $K_i = 39.8 \mu\text{M}$ ) with a potency similar to that of unlabeled AMP (Fig. 3C). On the other hand, while unlabeled AMP was able to block a large proportion of the binding of 10  $\mu\text{M}$   $^3\text{H}$ -adenosine to transfectants ( $K_i = 24.6 \mu\text{M}$ ), it could not block the binding as completely as unlabeled adenosine and had little effect in blocking  $^3\text{H}$ -adenosine binding to nontransfected HEK293 cells (Fig. 3D). While these results give support to the idea that both AMP and adenosine bind to P2Y15, because neither AMP nor adenosine can antagonize the binding of  $^3\text{H}$ -AMP to nontransfected HEK293 cells, the results also demonstrate a lack of specific AMP binding sites on the nontransfected cells.

## 20 EXAMPLE 20

To test whether any of the known non-specific antagonists of adenosine receptors could antagonize P2Y15, calcium mobilization assays was performed in the presence of varying concentrations of theophylline, 8-phenyltheophylline, 3-isobutyl-1 methylxanthine (IBMX), and caffeine. All of these compounds were able to block the calcium mobilization induced by AMP and adenosine, with  $K_i$  values for blocking AMP ranging from 250 nM for 8-phenyltheophylline to 2700 nM for caffeine and  $K_i$  values for blocking adenosine ranging from 260 nM for 8-phenyltheophylline to 26200 nM for caffeine. (Fig. 4A and B).

**EXAMPLE 21**

Adenosine 5'-( $\alpha,\beta$ -methylene)diphosphate (AMP-CP), which as a potent inhibitor of ectonucleotidases has been used in the past to provide evidence that blocking the conversion of AMP to adenosine can effectively block cellular responses to AMP, was effective in blocking calcium responses to not only AMP but also to adenosine, indicating that this compound can also directly inhibit binding to the P2Y<sub>15</sub> receptor (Fig. 4).

**EXAMPLE 22**

*Mast cell expression of P2Y<sub>15</sub> detected by microarray analysis*

*Target preparation*

Total RNA was prepared from human umbilical cord blood-derived mast cells (HCMC) according to the method for generating HCMC in vitro established by H. Saito (H. Saito et al. *J. Immunol.* 157:343-350, 1996). The HCMC so generated appear to resemble human lung mast cells in terms of their intracellular protease profiles, their histamine release characteristics, and their pharmacological properties (H. Saito et al. *J. Immunol.* 157:343-350;1996, Y. Igarashi et al. *Clin. Exp. Allergy* 26:597-602;1996, and H. Nagai et al. *Clin. Exp. Allergy* 28:1228-1236;1998). The HCMC, therefore, are useful for biological and pharmacological studies of lung mast cells and for the development of new anti-asthma drugs. Total RNA from the HCMC was isolated using Trizol<sup>TM</sup> (Invitrogen Corp., Carlsbad, CA, USA) according to the manufacturer's protocol. Five micrograms of the total RNA was then added to a reaction mix in a final volume of 12  $\mu$ l, containing bacterial control mRNAs (2.5 pg/ $\mu$ l araB/entF, 8.33 pg/ $\mu$ l fixB/gnd and 25 pg/ $\mu$ l hisB/leuB) and 1.0  $\mu$ l of 0.5 pmol/ $\mu$ l T7-(dT)<sub>24</sub> oligonucleotide primer. The mixture was incubated for 10 min at 70°C and chilled on ice. With the mixture remaining on ice, 4  $\mu$ l of 5x first-strand buffer, 2  $\mu$ l 0.1 M DTT, 1  $\mu$ l of 10 mM dNTP mix and 1  $\mu$ l Superscript<sup>TM</sup> II RNase H- reverse transcriptase (200 U/ $\mu$ l) was added to make a final volume of 20  $\mu$ l, and

the mixture incubated for 1 h in a 42°C water bath. Second-strand cDNA was synthesized in a final volume of 150 µl, in a mixture containing 30 µl of 5x second-strand buffer, 3 µl of 10 mM dNTP mix, 4 µl of *Escherichia coli* DNA polymerase I (10 U/µl) and 1 µl of RNase H (2 U/µl) for 2 h at 16°C. The cDNA was purified using a Qiagen QIAquick purification kit, dried down, and resuspended in IVT reaction mix, containing 3.0 µl nuclease-free water, 4.0 µl 10x reaction buffer, 4.0 µl 75 mM ATP, 4.0 µl 75 mM GTP, 3.0 µl 75 mM CTP, 3.0 µl 75 mM UTP, 7.5 µl 10 mM Biotin 11-CTP, 7.5 µl 10 mM Biotin 11-UTP (PerkinElmer Life Sciences Inc. Boston, MA, USA) and 4.0 µl enzyme mix. The reaction mix was incubated for 14 h at 37 °C and cRNA target purified using an RNeasy® kit (Qiagen). cRNA yield was quantified by measuring the UV absorbance at 260 nm, and fragmented in 40 mM Tris-acetate (TrisOAc) pH 7.9, 100 mM KOAc and 31.5 mM MgOAc, at 94°C for 20 min. This results typically in a fragmented target with a size range between 100 and 200 bases.

#### *Array hybridization*

Ten micrograms of fragmented target cRNA per array was used for hybridization to UniSet Human I and II Bioarrays (AmershamBiosciences) in 260 µl of hybridization solution containing 78 µl Amersham Hyb buffer component A and 130 µl Amersham Hyb buffer component B. The hybridization solution was heated at 90°C for 5 min to denature the cRNA and chilled on ice. The sample was vortexed for 5 s at maximum speed, and 250 µl injected into the inlet port of the hybridization chamber. The slides were loaded into a ISF-4-W shaking incubator (Kuhner, Birsfelden, Switzerland), with the hybridization chambers facing up. Slides were incubated for 24 h at 37°C, while shaking at 300 r.p.m.

#### *Post-hybridization processing using streptavidin-Cy5*

The slides were removed from the ISF-4-W shaker, and the hybridization chamber removed from each slide. Each slide was briefly rinsed in TNT buffer (0.1 M Tris-

HCl pH 7.6, 0.15 M NaCl, 0.05% Tween-20) at room temperature, and then washed in TNT buffer at 42 °C for 60 min. The signal was developed using a 1:500 dilution of streptavidin-Cy5 (AmershamBiosciences), for 30 min at room temperature. Excess dye was removed by washing four times with TNT buffer, for 5 min each, at room temperature. Slides were rinsed in 0.05% Tween-20 and dried under nitrogen gas. Processed slides were scanned using an Axon GenePix 4000B Scanner with the laser set to 635 nm, the photomultiplier tube (PMT) voltage to 600 and the scan resolution to 10µm. Images were acquired with the Axon GenePixPro v4.0 Scanning Software (AmershamBiosciences), and analyzed using the CodeLink™ Expression Analysis Software (AmershamBiosciences).

#### *Data analysis*

CodeLink™ Expression Analysis Software(AmershamBiosciences) automatically creates signal data for each spotted dot as a Microsoft Excel formatted spreadsheet. The data was analyzed using the computer program Spotfire Decision Site 7.0 (Spotfire Japan K.K., Tokyo, Japan) to determine the relative intensity of expression for each of the approximately 20,000 genes represented on the two arrays. As a result of this analysis, the P2Y15 gene transcript was found to be consistently expressed higher among the top approximately 1% of genes in terms of relative intensity. Fig. 5 shows the expression intensity of P2Y15 in relation to other well known genes expressed in mast cells.

### References

1. Nandanan E, Jang SY, Moro S, Kim HO, Siddiqui MA, Russ P, Marquez VE, Busson R, Herdewijn  
5
2. P, Harden TK, Boyer JL, Jacobson KA. Synthesis, biological activity, and molecular modeling of ribose-modified deoxyadenosine bisphosphate analogues as P2Y(1) receptor ligands. *J Med Chem.* 2000 Mar 9;43(5):829-42.  
10
3. Major DT, Halbfinger E, Fischer B. Molecular recognition of modified adenine nucleotides by the P2Y(1)-receptor. 2. A computational approach. *J Med Chem.* 1999 Dec 30;42(26):5338-47.  
15
3. Halbfinger E, Major DT, Ritzmann M, Uhl J, Reiser G, Boyer JL, Harden KT, Fischer B. Molecular recognition of modified adenine nucleotides by the P2Y(1)-receptor. 1. A synthetic, biochemical, and NMR approach. *J Med Chem.* 1999 Dec 30;42(26):5325-37.  
20
4. Moro S, Hoffmann C, Jacobson KA. Role of the extracellular loops of G protein-coupled receptors in ligand recognition: a molecular modeling study of the human P2Y1 receptor. *Biochemistry.* 1999 Mar 23;38(12):3498-507.  
25
5. Moro S, Guo D, Camaioni E, Boyer JL, Harden TK, Jacobson KA. Human P2Y1 receptor: molecular modeling and site-directed mutagenesis as tools to identify agonist and antagonist recognition sites. *J Med Chem.* 1998 Apr 23;41(9):1456-66.

6. Webb TE, Simon J, Barnard EA. Regional distribution of [35S]2'-deoxy 5'-O-(1-thio) ATP binding sites and the P2Y1 messenger RNA within the chick brain. *Neuroscience*. 1998 Jun;84(3):825-37.
- 5 7. Homolya L, Watt WC, Lazarowski ER, Koller BH, Boucher RC. Nucleotide-regulated calcium signaling in lung fibroblasts and epithelial cells from normal and P2Y(2) receptor (-/-) mice. *J Biol Chem*. 1999 Sep 10;274(37):26454-60.
- 10 8. Leon C, Hechler B, Vial C, Leray C, Cazenave JP, Gachet C. The P2Y1 receptor is an AMP OR ADENOSINE RECEPTOR LIGAND receptor antagonized by ATP and expressed in platelets and megakaryoblastic cells. *FEBS Lett*. 1997 Feb 10;403(1):26-30.
- 15 9. Dehaye JP, Moran A, Marino A. Purines, a new class of agonists in salivary glands? *Arch Oral Biol*. 1999 May;44 Suppl 1:S39-43.
- 20 10. King BF, Townsend-Nicholson A, Burnstock G. Metabotropic receptors for ATP and UTP: exploring the correspondence between native and recombinant nucleotide receptors. *Trends Pharmacol Sci*. 1998 Dec;19(12):506-14.
- 25 11. Saleh A, Figarella C, Kammouni W, Marchand-Pinatel S, Lazdunski A, Tubul A, Brun P, Merten MD. Pseudomonas aeruginosa quorum-sensing signal molecule N-(3-oxododecanoyl)-L-homoserine lactone inhibits expression of P2Y receptors in cystic fibrosis tracheal gland cells. *Infect Immun*. 1999 Oct;67(10):5076-82.
- 30 12. Cressman VL, Lazarowski E, Homolya L, Boucher RC, Koller BH, Grubb BR. Effect of loss of P2Y(2) receptor gene expression on nucleotide regulation of murine epithelial Cl(-) transport. *J Biol Chem*. 1999 Sep 10;274(37):26461-8.



Claims

1. A method for detecting the activity of P2Y15 in a sample comprising the steps of:
- 5
- a) incubating a sample comprising P2Y15 and a ligand under conditions which allow binding of P2Y15 and the ligand, and
  - b) detecting a second messenger, wherein said ligand is AMP or adenosine receptor ligand.
- 10
2. The method of claim 1 further comprising the steps of:
- a) incubating a second sample comprising P2Y15 in the absence of the ligand under conditions which allow binding of P2Y15 and the ligand, and
  - b) detecting a second messenger.
- 15
3. The method of claim 1 wherein said sample comprises cells expressing P2Y15.
- 20
4. The method of claim 1 wherein said sample comprises cell membranes bearing P2Y15.
5. A method of screening for an agent to modulate P2Y15 activity using cells expressing P2Y15, said method comprising:
- 25
- a) incubating a first sample of said cells in the presence of said agent and a second sample of said cells in the absence of said agent, both said samples under conditions which allow binding of AMP or adenosine receptor ligand to P2Y15;
- 0

- b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and
- c) comparing the results of said second messenger assays for said first and second samples.
- 5
6. A method of screening for an agent to modulate P2Y15 activity using cell membranes bearing P2Y15, said method comprising:
- a) incubating a first sample of said cell membranes in the presence of said agent and a second sample of said cell membranes in the absence of said agent, both said samples under conditions which allow binding of AMP or adenosine receptor ligand to P2Y15;
- 10 b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and
- 15 c) comparing the results of said second messenger assays for said first and second samples.
7. A method for determining if a test compound increases or decreases the activity of P2Y15 using cells expressing P2Y15, said method comprising:
- 20 a) incubating a first sample of said cells in the presence of said test compound and a second sample of said cells in the absence of said test compound, both said samples under conditions which permit binding of AMP or adenosine receptor ligand to P2Y15;
- 25 b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and
- c) comparing the results of said second messenger assays for said first and second samples.

- 90 -

8. A method for determining if a test compound increases or decreases the activity of P2Y15 using cell membranes bearing P2Y15, said method comprising:
- 5 a) incubating a first sample of said cell membranes in the presence of said test compound and a second sample of said cell membranes in the absence of said test compound, both said samples under conditions which permit binding of AMP or adenosine receptor ligand to P2Y15;
- 10 b) detecting a signalling activity of P2Y15 polypeptide in said first and second samples, and
- c) comparing the results of said second messenger assays for said first and second samples.
9. A method of identifying an agent that modulates the function of P2Y15, said method comprising:
- 15 a) contacting a P2Y15 polypeptide in the presence and absence of an agent under conditions permitting the binding of said AMP or adenosine receptor ligand to said P2Y15 polypeptide; and
- 20 b) measuring the binding of said P2Y15 polypeptide to said agent, relative to the binding in the absence of said agent, wherein an agent which changes binding is identified as a potential therapeutic agent for decreasing or increasing the function of P2Y15.
- 25 10. The method of claim 9 wherein said measuring is performed using a method selected from label displacement, surface plasmon resonance, fluorescence resonance energy transfer, fluorescence quenching, and fluorescence polarization.
- 30

11. The method of any one of claims 5 to 10 wherein said agent is selected from the group consisting of a natural or synthetic peptide, a polypeptide, an antibody or antigen-binding fragment thereof, a lipid, a carbohydrate, a nucleic acid, and a small organic molecule.
- 5 12. The method of any one of claims 5 to 10 wherein said step of measuring a signalling activity of said P2Y15 polypeptide comprises detecting a change in the level of a second messenger.
- 10 13. The method of either of claims 5 to 10 wherein the step of detecting a signalling activity comprises measurement of guanine nucleotide binding or exchange, adenylate cyclase activity, cAMP, protein kinase C activity, phosphatidylinositol breakdown, diacylglycerol, inositol triphosphate, intracellular calcium, arachinoid acid concentration, MAP kinase activity, tyrosine kinase activity, reporter gene expression.
- 15 14. A reagent that modulates the activity of a P2Y15 polypeptide or polynucleotide, wherein said reagent is identified by the method of any of the claims 1 to 13.
- 20 15. A pharmaceutical composition, comprising:  
the reagent of claim 14, and a pharmaceutically acceptable carrier.
- 25 16. Use of the reagent of claim 14 in the preparation of a medicament for modulating the activity of P2Y15 in a disease.
- 30 17. Use of claim 16, wherein the disease is a disease related to kidney function, disease related to mast cells, upper respiratory and oral disease, or allergic or inflammatory disease.

- 92 -

18. Use of claim 16, wherein the disease is one selected from the group consisting of asthma, COPD, congestive heart failure, high blood pressure, edema, cirrhosis of the liver, nephrotic syndrome, mast cell hyperplasia in the skin (mastocytosis), urticaria, flushing, single mastocytomas, multiple cutaneous lesions of urticaria pigmentosa, inflammatory bowel diseases, parasitic infections, cutaneous neurofibromas, benign and malignant breast lesions, urge incontinence, overactive bladder, Sjogren's syndrome, dry mouth, dental carries, post-nasal drip, and cough.
- 5

- 1 - JC17 Rec'd PCT/PTO 25 MAR 2005

## SEQUENCE LISTING

&lt;110&gt; Bayer Yakuhin, Ltd.

&lt;120&gt; REGULATION OF HUMAN P2Y15 G PROTEIN-COUPLED RECEPTOR

&lt;130&gt; x

&lt;160&gt; 6

&lt;170&gt; PatentIn version 3.2

&lt;210&gt; 1

&lt;211&gt; 1014

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1) .. (1014)

&lt;400&gt; 1

atg aat gag cca cta gac tat tta gca aat gct tot gat ttc ccc gat	48
Met Asn Glu Pro Leu Asp Tyr Leu Ala Asn Ala Ser Asp Phe Pro Asp	
1 5 10 15	
tat gca gct gct ttt gga aat tgc act gat gaa aac atc cca ctc aag	96
Tyr Ala Ala Phe Gly Asn Cys Thr Asp Glu Asn Ile Pro Leu Lys	
20 25 30	
atg cac tac ctc cct gtt att tat ggc att atc ttc ctc gtg gga ttt	144
Met His Tyr Leu Pro Val Ile Tyr Gly Ile Ile Phe Leu Val Gly Phe	
35 40 45	
cca ggc aat gca gta gtg ata tcc act tac att ttc aaa atg aga cct	192
Pro Gly Asn Ala Val Val Ile Ser Thr Tyr Ile Phe Lys Met Arg Pro	
50 55 60	
tgg aag agc agc acc atc att atg ctg aac ctg gcc tgc aca gat ctg	240
Trp Lys Ser Ser Thr Ile Ile Met Leu Asn Leu Ala Cys Thr Asp Leu	
65 70 75 80	
ctg tat ctg acc agc ctc ccc ttc ctg att cac tac tat gcc agt ggc	288
Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly	
85 90 95	
gaa aac tgg atc ttt gga gat ttc atg tgt aag ttt atc cgc ttc agc	336
Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Ser	
100 105 110	
ttc cat ttc aac ctg tat agc agc atc ctc ttc ctc acc tgt ttc agc	384
Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser	
115 120 125	
atc ttc cgc tac tgt gtg atc att cac cca atg agc tgc ttt toc att	432
Ile Phe Arg Tyr Cys Val Ile Ile His Pro Met Ser Cys Phe Ser Ile	
130 135 140	
cac aaa act cga tgt gca gtt gta gcc tgt gct gtg gtg tgg atc att	480
His Lys Thr Arg Cys Ala Val Val Ala Cys Ala Val Val Trp Ile Ile	
145 150 155 160	

- 2 -

tca ctg gta gct gtc att ccg atg acc ttc ttg atc aca tca acc aac	528
Ser Leu Val Ala Val Ile Pro Met Thr Phe Leu Ile Thr Ser Thr Asn	
165 170 175	
agg acc aac aga tca gcc tgt ctc gac ctc acc agt tcg gat gaa ctc	576
Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Glu Leu	
180 185 190	
aat act att aag tgg tac aac ctg att ttg act gca act act ttc tgc	624
Asn Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys	
195 200 205	
ctc ccc ttg gtg ata gtg aca ctt tgc tat acc acg att atc cac act	672
Leu Pro Leu Val Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile His Thr	
210 215 220	
ctg acc cat gga ctg caa act gac agc tgc ctt aag cag aaa gca cga	720
Leu Thr His Gly Leu Gln Thr Asp Ser Cys Leu Lys Gln Lys Ala Arg	
225 230 235 240	
agg cta acc att ctg cta ctc ctt gca ttt tac gta tgt ttt tta ccc	768
Arg Leu Thr Ile Leu Leu Leu Ala Phe Tyr Val Cys Phe Leu Pro	
245 250 255	
ttc cat atc ttg agg gtc att cgg atc gaa tct cgc ctg ctt tca atc	816
Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile	
260 265 270	
agt tgt tcc att gag aat cag atc cat gaa gct tac atc gtt tct aga	864
Ser Cys Ser Ile Glu Asn Gln Ile His Glu Ala Tyr Ile Val Ser Arg	
275 280 285	
cca tta gct gct ctg aac acc ttt ggt aac ctg tta cta tat gtg gtg	912
Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Leu Tyr Val Val	
290 295 300	
gtc agc gac aac ttt cag cag gct gtc tgc tca aca gtg aga tgc aaa	960
Val Ser Asp Asn Phe Gln Gln Ala Val Cys Ser Thr Val Arg Cys Lys	
305 310 315 320	
gta agc ggg aac ctt gag caa gca aag aaa att agt tac tca aac aac	1008
Val Ser Gly Asn Leu Glu Gln Ala Lys Lys Ile Ser Tyr Ser Asn Asn	
325 330 335	
cct tga	1014
Pro	

<210> 2  
 <211> 337  
 <212> PRT  
 <213> Homo sapiens

<400> 2

Met Asn Glu Pro Leu Asp Tyr Leu Ala Asn Ala Ser Asp Phe Pro Asp
1 5 10 15

Tyr Ala Ala Ala Phe Gly Asn Cys Thr Asp Glu Asn Ile Pro Leu Lys
20 25 30

- 3 -

Met His Tyr Leu Pro Val Ile Tyr Gly Ile Ile Phe Leu Val Gly Phe  
 35 40 45

Pro Gly Asn Ala Val Val Ile Ser Thr Tyr Ile Phe Lys Met Arg Pro  
 50 55 60

Trp Lys Ser Ser Thr Ile Ile Met Leu Asn Leu Ala Cys Thr Asp Leu  
 65 70 75 80

Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly  
 85 90 95

Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Ser  
 100 105 110

Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser  
 115 120 125

Ile Phe Arg Tyr Cys Val Ile Ile His Pro Met Ser Cys Phe Ser Ile  
 130 135 140

His Lys Thr Arg Cys Ala Val Val Ala Cys Ala Val Val Trp Ile Ile  
 145 150 155 160

Ser Leu Val Ala Val Ile Pro Met Thr Phe Leu Ile Thr Ser Thr Asn  
 165 170 175

Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Glu Leu  
 180 185 190

Asn Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys  
 195 200 205

Leu Pro Leu Val Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile His Thr  
 210 215 220

Leu Thr His Gly Leu Gln Thr Asp Ser Cys Leu Lys Gln Lys Ala Arg  
 225 230 235 240

Arg Leu Thr Ile Leu Leu Leu Leu Ala Phe Tyr Val Cys Phe Leu Pro  
 245 250 255

Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile  
 260 265 270

Ser Cys Ser Ile Glu Asn Gln Ile His Glu Ala Tyr Ile Val Ser Arg  
 275 280 285



- 4 -

Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Leu Tyr Val Val  
 290 295 300

Val Ser Asp Asn Phe Gln Gln Ala Val Cys Ser Thr Val Arg Cys Lys  
 305 310 315 320

Val Ser Gly Asn Leu Glu Gln Ala Lys Lys Ile Ser Tyr Ser Asn Asn  
 325 330 335

Pro

<210> 3  
 <211> 1014  
 <212> DNA  
 <213> Mus musculus

<220>  
 <221> CDS  
 <222> (1)..(1014)

<400> 3  
 atg att gag cca ctg gac agt cca gcc agt gat tgg gat ttc ctg gat 48  
 Met Ile Glu Pro Leu Asp Ser Pro Ala Ser Asp Ser Asp Phe Leu Asp  
 1 5 10 15  
 tac cca agt gct ctg gga aac tgc acc gac gag caa atc tca ttc aag 96  
 Tyr Pro Ser Ala Leu Gly Asn Cys Thr Asp Glu Gln Ile Ser Phe Lys  
 20 25 30  
 atg cag tac ctt ccc gtc atc tat agc atc atc ttc ctc gtg ggc ttc 144  
 Met Gln Tyr Leu Pro Val Ile Tyr Ser Ile Ile Phe Leu Val Gly Phe  
 35 40 45  
 ccg ggg aac aca gtg gcc atc tcc atc tac att ttc aag atg cgg ccg 192  
 Pro Gly Asn Thr Val Ala Ile Ser Ile Tyr Ile Phe Lys Met Arg Pro  
 50 55 60  
 tgg agg ggc agt aca gtc atc atg ctg aac ctg gcc ttg acg gac ttg 240  
 Trp Arg Gly Ser Thr Val Ile Met Leu Asn Leu Ala Leu Thr Asp Leu  
 65 70 75 80  
 ctg tat ctg acc agc ctc ccg ttc ctc atc cat tac tat gcc agt ggt 288  
 Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly  
 85 90 95  
 gaa aac tgg atc ttt gga gat ttc atg tgc aag ttc atc cgc ttc ggc 336  
 Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Gly  
 100 105 110  
 ttc cac ttc aac ctc tac agc agc att ctc ttc ctc acc tgc ttc agt 384  
 Phe His Phe Asn Leu Tyr Ser Ile Leu Phe Leu Thr Cys Phe Ser  
 115 120 125  
 ctc ttc cgt tac gtt gtg atc att cac ccg atg agc tgc ttt tct att 432  
 Leu Phe Arg Tyr Val Val Ile Ile His Pro Met Ser Cys Phe Ser Ile  
 130 135 140

- 5 -

cag aaa act cgc tgg gca gtg gta gct tgt gcc ggg gtg tgg gtc att 480  
 Gln Lys Thr Arg Trp Ala Val Val Ala Cys Ala Gly Val Trp Val Ile  
 145 150 155 160

tot ttg gta gct gtc atg ccc atg act ttc ctg atc aca tca acc acc 528  
 Ser Leu Val Ala Val Met Pro Met Thr Phe Leu Ile Thr Ser Thr Thr  
 165 170 175

cgg acc aat agg tct gct tgc ctt gac ctc acc agt tca gat gac ctc 576  
 Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Asp Leu  
 180 185 190

act act atc aag tgg tac aat ctc att ttg aca gcc acc act ttc tgc 624  
 Thr Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys  
 195 200 205

ctg cca ttg gtg ata gtg aca ctt tgc tac acg aca att atc agt acc 672  
 Leu Pro Leu Val Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile Ser Thr  
 210 215 220

ctg act cac ggg cct cgg acc cac agc tgc ttt aag cag aag gct cgg 720  
 Leu Thr His Gly Pro Arg Thr His Ser Cys Phe Lys Gln Lys Ala Arg  
 225 230 235 240

aga ctg act att ctg ctc ctc ctt gtt ttc tat ata tgt ttc tta ccc 768  
 Arg Leu Thr Ile Leu Leu Leu Leu Val Phe Tyr Ile Cys Phe Leu Pro  
 245 250 255

ttc cac atc ttg agg gtc att cgg atc gaa tct cgc ctg ctt tca atc 816  
 Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile  
 260 265 270

agc tgc tcc atc gag agt cac atc cac gaa gct tac att gtt tct aga 864  
 Ser Cys Ser Ile Glu Ser His Ile His Glu Ala Tyr Ile Val Ser Arg  
 275 280 285

cca tta gct gct ctc aac acc ttt ggc aac ctg ctg tta tat gtt gtg 912  
 Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Tyr Val Val  
 290 295 300

gtc agc aat aac ttc cag cag gca ttc tgc tot ata gtg aga tgc aaa 960  
 Val Ser Asn Asn Phe Gln Gln Ala Phe Cys Ser Ile Val Arg Cys Lys  
 305 310 315 320

gcc agt ggg gac ctt gaa caa gga aag aaa gac agt tgc tca aac aac 1008  
 Ala Ser Gly Asp Leu Glu Gln Gly Lys Lys Asp Ser Cys Ser Asn Asn  
 325 330 335

cct tga 1014  
 Pro

<210> 4  
 <211> 337  
 <212> PRT  
 <213> Mus musculus

<400> 4

Met Ile Glu Pro Leu Asp Ser Pro Ala Ser Asp Ser Asp Phe Leu Asp  
 1 5 10 15

- 6 -

Tyr Pro Ser Ala Leu Gly Asn Cys Thr Asp Glu Gln Ile Ser Phe Lys  
 20 25 30

Met Gln Tyr Leu Pro Val Ile Tyr Ser Ile Ile Phe Leu Val Gly Phe  
 35 40 45

Pro Gly Asn Thr Val Ala Ile Ser Ile Tyr Ile Phe Lys Met Arg Pro  
 50 55 60

Trp Arg Gly Ser Thr Val Ile Met Leu Asn Leu Ala Leu Thr Asp Leu  
 65 70 75 80

Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly  
 85 90 95

Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Gly  
 100 105 110

Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser  
 115 120 125

Leu Phe Arg Tyr Val Val Ile Ile His Pro Met Ser Cys Phe Ser Ile  
 130 135 140

Gln Lys Thr Arg Trp Ala Val Val Ala Cys Ala Gly Val Trp Val Ile  
 145 150 155 160

Ser Leu Val Ala Val Met Pro Met Thr Phe Leu Ile Thr Ser Thr Thr  
 165 170 175

Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Asp Leu  
 180 185 190

Thr Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys  
 195 200 205

Leu Pro Leu Val Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile Ser Thr  
 210 215 220

Leu Thr His Gly Pro Arg Thr His Ser Cys Phe Lys Gln Lys Ala Arg  
 225 230 235 240

Arg Leu Thr Ile Leu Leu Leu Val Phe Tyr Ile Cys Phe Leu Pro  
 245 250 255

Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile  
 260 265 270

- 7 -

Ser Cys Ser Ile Glu Ser His Ile His Glu Ala Tyr Ile Val Ser Arg  
275 280 285

Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Tyr Val Val  
290 295 300

Val Ser Asn Asn Phe Gln Gln Ala Phe Cys Ser Ile Val Arg Cys Lys  
305 310 315 320

Ala Ser Gly Asp Leu Glu Gln Gly Lys Lys Asp Ser Cys Ser Asn Asn  
325 330 335

Pro

<210> 5  
<211> 1014  
<212> DNA  
<213> Rattus norvegicus

<220>  
<221> CDS  
<222> (1)..(1014)

<400> 5  
atg att gag aca ctg gac agc cca gcc aat gat tct gat ttc ctg gat 48  
Met Ile Glu Thr Leu Asp Ser Pro Ala Asn Asp Ser Asp Phe Leu Asp  
1 5 10 15  
tac ata act gct ttg gaa aac tgc act gat gag caa atc tca ttc aag 96  
Tyr Ile Thr Ala Leu Glu Asn Cys Thr Asp Glu Gln Ile Ser Phe Lys  
20 25 30  
atg cag tac ctt ccc gtc atc tac agc atc atc ttt otc gtg ggc ttc 144  
Met Gln Tyr Leu Pro Val Ile Tyr Ser Ile Ile Phe Leu Val Gly Phe  
35 40 45  
cag gga aat acg gtg gcg att tcc atc tac gtt ttc aag atg cga cct 192  
Pro Gly Asn Thr Val Ala Ile Ser Ile Tyr Val Phe Lys Met Arg Pro  
50 55 60  
tgg aag agc agt acc atc atc atg ctg aac ctg gcc ttg acg gac ttg 240  
Trp Lys Ser Ser Thr Ile Ile Met Leu Asn Leu Ala Leu Thr Asp Leu  
65 70 75 80  
ctg tat ctg acc ago otc cct ttc ctc atc cat tat tac gcg agc ggt 288  
Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly  
85 90 95  
gaa aac tgg atc ttc ggg gat ttc atg tgc aag ttc atc cga ttt ggc 336  
Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Gly  
100 105 110  
ttc cat ttc aac ctt tac agc agc atc otc ttc otc acc tgc ttt agc 384  
Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser  
115 120 125

- 8 -

ctc ttc cgt tac att gtg atc att cac ccg atg agc tgt ttt tct att Leu Phe Arg Tyr Ile Val Ile Ile His Pro Met Ser Cys Phe Ser Ile 130 135 140	432
cag aag act cga tgg gcg gtg gtg gct tgt gct ggg gtg tgg gtc att Gln Lys Thr Arg Trp Ala Val Val Ala Cys Ala Gly Val Trp Val Ile 145 150 155 160	480
tct ttg gta gct gtc atg ccc atg act ttc ctg atc aca tca acc acc Ser Leu Val Ala Val Met Pro Met Thr Phe Leu Ile Thr Ser Thr Thr 165 170 175	528
cgg acc aat agg tct gct tgc ctt gac ctc acc agc tca gat gac ctc Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Asp Leu 180 185 190	576
act act atc aaa tgg tac aat ctc att ttg acg gct acc act ttc tgc Thr Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys 195 200 205	624
ctg ccc ttg ctg ata gtg aca ctc tgc tac acg acg att atc agc acc Leu Pro Leu Leu Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile Ser Thr 210 215 220	672
ctg act cac gga cct cgg acc cac agc tgc ttt aag cag aag gct cgg Leu Thr His Gly Pro Arg Thr His Ser Cys Phe Lys Gln Lys Ala Arg 225 230 235 240	720
agg ctg acg atc ctg ctc ctc ctt gtg ttc tat gta tgc ttt tta ccc Arg Leu Thr Ile Leu Leu Leu Val Phe Tyr Val Cys Phe Leu Pro 245 250 255	768
ttc cac atc ctt agg gtc att cgg atc gaa tct cgc ctg ctt tca atc Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile 260 265 270	816
agc tgc tcc atc gag agt cac atc cac gaa gct tac att gtc tct agg Ser Cys Ser Ile Glu Ser His Ile His Glu Ala Tyr Ile Val Ser Arg 275 280 285	864
cca tta gct gcc ctc aac acc ttt ggc aac ctg ctg tta tat gtc gtc Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Leu Tyr Val Val 290 295 300	912
gtc agc aat aac ttc cag cag gca ttc tgc tcc gca gtg aga tgt aaa Val Ser Asn Asn Phe Gln Gln Ala Phe Cys Ser Ala Val Arg Cys Lys 305 310 315 320	960
gcc atc ggg gac ctt gaa caa gca aag aaa gac agt tgc tca aac aac Ala Ile Gly Asp Leu Glu Gln Ala Lys Lys Asp Ser Cys Ser Asn Asn 325 330 335	1008
ccc tga Pro	1014

<210> 6  
 <211> 337  
 <212> PRT  
 <213> Rattus norvegicus

- 9 -

&lt;400&gt; 6

Met Ile Glu Thr Leu Asp Ser Pro Ala Asn Asp Ser Asp Phe Leu Asp  
 1 5 10 15

Tyr Ile Thr Ala Leu Glu Asn Cys Thr Asp Glu Gln Ile Ser Phe Lys  
 20 25 30

Met Gln Tyr Leu Pro Val Ile Tyr Ser Ile Ile Phe Leu Val Gly Phe  
 35 40 45

Pro Gly Asn Thr Val Ala Ile Ser Ile Tyr Val Phe Lys Met Arg Pro  
 50 55 60

Trp Lys Ser Ser Thr Ile Ile Met Leu Asn Leu Ala Leu Thr Asp Leu  
 65 70 75 80

Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly  
 85 90 95

Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Gly  
 100 105 110

Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser  
 115 120 125

Leu Phe Arg Tyr Ile Val Ile Ile His Pro Met Ser Cys Phe Ser Ile  
 130 135 140

Gln Lys Thr Arg Trp Ala Val Val Ala Cys Ala Gly Val Trp Val Ile  
 145 150 155 160

Ser Leu Val Ala Val Met Pro Met Thr Phe Leu Ile Thr Ser Thr Thr  
 165 170 175

Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Asp Leu  
 180 185 190

Thr Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys  
 195 200 205

Leu Pro Leu Leu Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile Ser Thr  
 210 215 220

Leu Thr His Gly Pro Arg Thr His Ser Cys Phe Lys Gln Lys Ala Arg  
 225 230 235 240

Arg Leu Thr Ile Leu Leu Leu Leu Val Phe Tyr Val Cys Phe Leu Pro  
 245 250 255

- 10 -

Phe His Ile Leu Arg Val Ile Arg Ile Glu Ser Arg Leu Leu Ser Ile  
260 265 270

Ser Cys Ser Ile Glu Ser His Ile His Glu Ala Tyr Ile Val Ser Arg  
275 280 285

Pro Leu Ala Ala Leu Asn Thr Phe Gly Asn Leu Leu Leu Tyr Val Val  
290 295 300

Val Ser Asn Asn Phe Gln Gln Ala Phe Cys Ser Ala Val Arg Cys Lys  
305 310 315 320

Ala Ile Gly Asp Leu Glu Gln Ala Lys Lys Asp Ser Cys Ser Asn Asn  
325 330 335

Pro

<210> 7  
<211> 22  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) .. (22)

<400> 7  
ttcggatcga atctcgctg ct

22

<210> 8  
<211> 23  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) .. (23)

<400> 8  
tgcttgctca aggttccgc tta

23

<210> 9  
<211> 28  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) .. (28)

<400> 9  
gccaaactga actctcttgt tttcttgc

28

- 11 -

<210> 10  
<211> 24  
<212> DNA  
<213> Homo sapiens


<220>  
<221> misc\_feature  
<222> (1)..(24)

<400> 10  
gccctggctt tggcacatga ttac



24





- 1/6 -

**Fig. 1**



- 2/6 -

**Fig. 2A****Fig. 2B**



- 3/6 -

**Fig. 3A****Fig. 3B**


- 4/6 -

**Fig. 3C****Fig. 3D**

- 5/6 -

**Fig. 4A****Fig. 4B**

- 6/6 -

**Fig. 5**

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 03/10666

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 G01N33/566 C07K14/705 C07K14/72		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 G01N C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, SEQUENCE SEARCH, WPI Data, PAJ, MEDLINE, EMBASE, BIOSIS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02 46414 A (SQUIBB BRISTOL MYERS CO) 13 June 2002 (2002-06-13) the whole document ---	1-18
X	WO 02 14511 A (BAYER AG ; RAMAKRISHNAN SHYAM (US)) 21 February 2002 (2002-02-21) cited in the application the whole document ---	1-18
X	WO 01 87980 A (APPLERA CORP) 22 November 2001 (2001-11-22) the whole document ---	1-18
X	WO 02 057441 A (MIYAJIMA NOBUYUKI ; ITO TAKASHI (JP); MIWA MASANORI (JP); SHINTANI) 25 July 2002 (2002-07-25) the whole document --- <div style="text-align: center;">-/-</div>	1-18
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents :</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center; font-weight: bold;">23 January 2004</div>		Date of mailing of the international search report  <div style="text-align: center; font-weight: bold;">11/02/2004</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center; font-weight: bold;">Pinheiro Vieira, E</div>

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 03/10666

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LEE D K ET AL: "Discovery and mapping of ten novel G protein-coupled receptor genes"</p> <p>GENE: AN INTERNATIONAL JOURNAL ON GENES AND GENOMES, ELSEVIER SCIENCE PUBLISHERS, BARKING, GB,</p> <p>vol. 275, no. 1,</p> <p>5 September 2001 (2001-09-05), pages 83-91, XP004307114</p> <p>ISSN: 0378-1119</p> <p>page 83 -page 91</p> <p>----</p>	1-18
P,X	<p>WO 02 088183 A (PARADIGM THERAPEUTICS LTD ;HENDRICK ALAN (GB); APARICIO SAMUEL (GB)</p> <p>7 November 2002 (2002-11-07)</p> <p>the whole document</p> <p>-----</p>	1-18



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP 03/10666

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0246414	A	13-06-2002	AU 3065102 A	18-06-2002
			WO 0246414 A2	13-06-2002
			US 2003157598 A1	21-08-2003
			US 2003224458 A1	04-12-2003
WO 0214511	A	21-02-2002	AU 9545401 A	25-02-2002
			WO 0214511 A2	21-02-2002
			EP 1311677 A2	21-05-2003
WO 0187980	A	22-11-2001	AU 6171501 A	26-11-2001
			CA 2409108 A1	22-11-2001
			EP 1287032 A2	05-03-2003
			WO 0187980 A2	22-11-2001
			US 2003049790 A1	13-03-2003
WO 02057441	A	25-07-2002	WO 02057441 A1	25-07-2002
			JP 2002355072 A	10-12-2002
WO 02088183	A	07-11-2002	WO 02088183 A2	07-11-2002

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**